

CIDETRAK® OFM MEC™

A Mating Disruption Formulation For:
oriental fruit moth, *Grapholita molesta*, macadamia nut borer, *Cryptophlebia ambrodelta*,
and koa seedworm, *Cryptophlebia illepidia*

ACTIVE INGREDIENT: **WT. BY %**
Z-8-Dodecen-1-yl Acetate.....9.05%
E-8-Dodecen-1-yl Acetate.....0.82%
Z-8-Dodecen-1-ol0.13%
OTHER INGREDIENTS:90.0%

TOTAL:100.00%

EPA Reg. No.: 51934-25 EPA Establishment No.: 51934-OK-001

NET CONTENTS:

..... [Fl. Oz./mL/Gals.]

MANUFACTURED BY: TRÉCÉ INCORPORATED, 7569 Hwy 28 W • Adair, Oklahoma 74330

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.

HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday through Friday, 8:00 a.m. to 12:00 p.m. pacific time or your poison control center at 1-800-222-1222. For additional information, you may also contact Trécé Inc. at 1-866-785-1313.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:	<ul style="list-style-type: none">• Long-sleeved shirt and long pants• Waterproof gloves• Shoes plus socks
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.	

USER SAFETY RECOMMENDATIONS

Users should:	<ul style="list-style-type: none">• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
----------------------	---

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

SHAKE WELL BEFORE USING

DO NOT FREEZE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For requirements specific to your state or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 0 hours.

CIDETRAK® OFM MEC™ is a suspension concentrate that disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation and reduces the need for applying insecticides.

Per 40 CFR 180.1153 DO NOT EXCEED the rate of 150 g A.I. per acre per year.

When using **CIDETRAK® OFM MEC™** in combination with other products, use **CIDETRAK® OFM MEC™** at the rate specified in **CIDETRAK® OFM MEC™** Rate Directions below. Follow the directions for use, precautions and limitations for use on all of the product labels used in the combination.

Crop	Peaches, Nectarines, Apricots, Plums, Cherries, and other stone fruits; Apples, Pears, Quince and other pome fruit; Almonds, Macadamia, and other nut crops; other crops where oriental fruit moth is a pest.
Targeted Pest	Oriental fruit moth (<i>Grapholita molesta</i>), macadamia nut borer (<i>Cryptophlebia ambrodelta</i>) and koa seedworm (<i>Cryptophlebia illepidia</i>)
Rate	Mix CIDETRAK® OFM MEC™ with prescribed and/or lower label rates of a suitable insecticide, fungicide, kairomone, oil and/or adjuvant. Also, may be applied to an orchard under current OFM pheromone based mating disruption products. A spreader sticker may be used for enhanced adhesion. Use CIDETRAK® OFM MEC™ at a rate 2.36-6.76 fl. oz/acre (70 to 200 mL/acre) (7.0-20.0 g A.I./acre) or 5.91-16.9 fl. oz/ha (175-500 mL/ha) (17.5-50.0 g A.I./ha). Apply CIDETRAK® OFM MEC™ usually at a rate of 2.36 fl. oz (70 mL) (7.0 g A.I.) per acre or 5.91 fl. oz (175 mL) (17.5 g A.I./ha) alone or in a mixture only with a suitable registered insecticide, fungicide, oil, and/or adjuvants
Ground/Aerial Application	Apply foregoing alone or in a mixture in a full coverage spray, according to directions for use of the insecticide, fungicide, kairomone, oil, and/or adjuvant. Coverage should be complete but do not allow "run-off."
Spray/Sprayer Preparation	Shake CIDETRAK® OFM MEC™ vigorously before measuring for spray solution. Fill sprayer half full of water. Start physical or by-pass agitation on spray application equipment. Carefully measure and pour or place insecticide, fungicide, kairomone, oil, and/or adjuvant into the spray tank. Carefully measure and pour the required amount of CIDETRAK® OFM MEC™ into the spray tank. Fill spray tank to the required level and initiate application.
Chemical Compatibility	Check physical compatibility of CIDETRAK® OFM MEC™ + insecticide, fungicide, kairomone, oil, and/or adjuvant before mixing in spray tanks. No plant phytotoxicity has been observed with mixtures of CIDETRAK® OFM MEC™ . But if in doubt, one may check plant phytotoxic response to mixtures of CIDETRAK® OFM MEC™ + insecticide, fungicide, kairomone, oil, and/or adjuvant by treating 4-5 plants or small area of foliar surface of plantings with chosen mixtures. Observe the effect over 2 days.
Monitoring/Timing	Apply CIDETRAK® OFM MEC™ prior to moth emergence or in between any generation. Retreatment may be necessary if there is an excessive amount of rain. Monitor moth activity by using PHEROCON® monitoring systems and lures for OFM. Begin monitoring before first emergence or when possible and continue throughout the season to assess population, phenology and abundance. Apply CIDETRAK® OFM MEC™ in mixtures according to the directions for timing of the manufacturer of the insecticide. Repeat applications of CIDETRAK® OFM MEC™ in mixtures according to directions for the insecticide.
Treatment Tips	Supplement CIDETRAK® OFM MEC™ using timely applications of insecticides to control high populations. Monitor all insect populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: 1. Treatment of external sources of infestation with CIDETRAK® OFM MEC™ 2. Treatment of external sources of infestation with an insecticide. 3. Treatment of pheromone treated orchard with insecticide. Consult your local extension specialist, certified crop consultant, or Trécé® representative for local field condition management strategies.

STORAGE AND DISPOSAL

Nonrefillable container. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store unopened original package in a dry location at a temperature below 40° F. Store in a cool, dry place, out of direct sunlight, and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in the original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

NONREFILLABLE CONTAINER (equal to or less than 5 gallons): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NONREFILLABLE CONTAINER [Greater Than 5 Gallons]: Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

WARRANTY AND LIMITATION OF DAMAGES

To the extent consistent with applicable law, all statements concerning the use of this product apply only when used as directed as stated on this product label. Trécé®, Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trécé®, Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.



® Mailing Address: P.O. Box 129 • Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W. • Adair, Oklahoma 74330
Phone: (918) 785-3061 • Fax: (918) 785-3063