

## **LIQUID BORON 10%**

GENERAL INFORMATION CM LIQUID BORON 10% is formulated for field, vegetable, row, fruit, tree, vine, ornamental and turf crops.

CM LIQUID BORON 10% may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume).

CM LIQUID BORON 10% may be applied alone or with fertilizers or pesticides. It may be applied either to the plant foliage or to the soil. CM LIQUID BORON 10% is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

**DIRECTIONS** 

Mixing Sequence

- 1. Water
- 2. CM LIQUID BORON 10%
- 3. Pesticide

Always add this product to water or nitrogen solution before adding phosphate material.

WARNING: CM LIQUID BORON 10% contains a percentage of Boron that can be detrimental to growing crops. Use of excessive amounts or on crops not recommended may result in serious injury to those crops. Excessive amounts of Boron may be toxic. Contact your local agricultural authority for additional information.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

## **GUARANTEED ANALYSIS**

Boron (B)......10.0%

Derived from Boric Acid

Wt/gal: 10.1 lb.@ 68°F Gr/ltr: 1209 g @ 20°C

RATES FOR USE

Foliar Application: Apply 2 to 4 quarts per acre as needed on a 10-14 day schedule. When rates of over 2 quarts are required, split applications of 2 quarts or less are recommended. Avoid foliar applications of CM LIQUID BORON 10% when plants are under stress due to lack of moisture.

Soil Application:

Maintenance: 1-3 qt/acre Mild deficiency: 2-6 qt/acre High deficiency: 1-3 gal/acre

Aerial Application: Apply 1-2 quarts per 5-30 gallons of water per acre. A concentration of 1 quart per gallon of spray may

be used.

Dilute Application: Apply the suggested rate of this product in enough water for proper coverage.

Concentrate Application: Use up to 2 quarts of this product in a minimum of 25 gallons of water per acre.

Drip Irrigation Application: Apply the suggested rate of this product in multiple applications 10-14 days apart throughout

the growing season of the crop.

## KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eve and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRSTAID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all governmental procedures for the disposal of pesticide spray solutions. Rinse container; empty rinsate into spray tank. Dispose of empty container according to approved local practices. Contact county agricultural commissioner for disposal information.

## CONDITIONS OF SALE

The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, the manufacturer does not make nor does it authorize any agent or representative to make any warranty or representation, expressed or implied, concerning this material or the use thereof except in conformity with the statements on the label. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions. If these conditions are unacceptable to the buyer, he should immediately return the unopened product to the 2-24

