
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1454A-S1 0222

FLUDIOXONIL	GROUP	12	FUNGICIDE
-------------	-------	-----------	-----------

Cannonball® WG

Fungicide

This supplemental label expires on 02/04/2025 and must not be used or distributed after this date.

Active Ingredient:	
Fludioxonil (CAS No. 131341-86-1)	50.0%
Other Ingredients:	50.0%
Total:	100.0%

Cannonball WG is a 50% water dispersible granule.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1454

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Cannonball WG as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Rotational Crops	Planting Time from Last Cannonball WG Application
Almond Beans (dried and succulent except cowpeas) Berries (bushberries 13-07B, caneberries 13-07A) Vegetable, Brassica, Head and Stem (Crop Group 5-16) Brassica, Leafy greens, except watercress (Subgroup 4-16B) Carrot Celtuce Cucurbit vegetables Crop Group 9 Fennel, Florence, fresh leaves and stalk Filbert Ginseng Herbs (fresh and dried) Kohlrabi Leafy Greens (Crop Subgroup 4-16A) Leaf petiole vegetables (Crop Subgroup 22B) Melons (Crop Subgroup 9A) Onion, Bulb, Crop Subgroup 3-07A; Onion, Green, Crop Subgroup 3-07B Fruiting vegetables Crop Group 8-10 Pecan Pistachio Tuberous and Corm Vegetables Subgroup 1C Leaves of Root and Tuber Vegetables Crop Group 2 Strawberries Tomatoes Vegetable, root, except sugar beet, subgroup 1B Walnut Watercress Crops Not Intended for Food or Feed	0 days
All Other Crops Intended for Food or Feed	30 days

CROP USE DIRECTIONS - FOR FOLIAR APPLIED

Crop	Disease	Rate oz/Acre	Use Directions
Leafy Greens, Crop Subgroup 4-16A, Lettuce, head and leaf Parsley Spinach Leaf Petiole Vegetables Crop Subgroup 22B, Celery Celtuce, Fennel, Florence, fresh leaves and stalk And cultivars and/or hybrids of these.	Gray mold <i>(Botrytis cinerea)</i> Sclerotinia rot <i>(Sclerotinia spp.)</i> Basal rot <i>(Phoma exigua)</i>	7*	<p>For foliar diseases, begin applications prior to or at the onset of disease and repeat applications on a 7-10 day interval if conditions remain favorable for disease development.</p> <p>For soil-borne disease:</p> <p>Direct Seeded lettuce: Apply immediately after emergence or prior to disease development.</p> <p>Transplanted lettuce: Apply immediately after transplanting or prior to disease development.</p> <p>A second application must be made if either, 1) the soil is disturbed by cultivation or thinning or, 2) conditions continue to favor disease. Apply no closer than a 7-day interval.</p> <p>Resistance Management: After 2 applications of Cannonball WG, alternate with another fungicide with a different mode of action for 2 applications.</p>

*7 oz product is equivalent to 0.22 lb ai fludioxonil/A

Complete List of Leafy Greens Crop Subgroup 4-16A: Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities

Leaf Petiole Vegetables Crop Subgroup 22B includes: Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. Make additional applications by ground or chemigation. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- 1) Maximum Single Application Rate: **DO NOT** exceed the maximum rate listed in the table.
- 2) **DO NOT** apply more than 4 applications at the maximum application rate per year.
- 3) Minimum Application Interval: 7 days
- 4) **DO NOT** make more than two applications by air.
- 5) **DO NOT** apply more than 28 oz/A (0.9 lb ai/A) of Cannonball WG per year.
- 6) **DO NOT** apply more than 0.9 lb ai/A of fludioxonil-containing products per plot of land per year.
- 7) May be applied on the day of harvest (0-day PHI).

Crop	Disease	Rate oz/100 gal	Use Directions
Cucumber (Greenhouse production only) (Not for use in California)	Fusarium Root Rot (<i>Fusarium solani</i>)	1.0 oz/100gal water (solution)*	Prepare a drench solution of 1 ounce Cannonball fungicide per 100 gallons water (solution). Apply 5-8 fl. oz. of solution onto the growing medium at the base of each cucumber plant.

Specific Use Restrictions:

- 1) Maximum Single Application Rate: **DO NOT** exceed the maximum rate listed in the table.
- 2) **DO NOT** apply more than 5-8 fl. oz. of solution by drip/drench application per treatment.
- 3) **DO NOT** apply more than 1 application per crop.
- 4) **DO NOT** apply to the foliage.
- 5) **DO NOT** apply within 1 day of harvest (1-day PHI).
- 6) **DO NOT** use in greenhouse for transplant production.

Crop	Disease	Rate oz/100 gal	Use Directions
Tomato (Greenhouse production only) (Not for use in California)	Fusarium Root Rot (Fusarium solani)	1.0 oz/100gal water (solution)*	Prepare a drench solution of 1 ounce Cannonball fungicide per 100 gallons water (solution). Apply 8 fl. oz. of solution onto the growing medium at the base of each tomato plant. If needed, make a second application 21 day after the first application.

*1 oz product is equivalent to 0.03 lb ai fludioxonil/100 gal of water

Specific Use Restrictions:

- 1) Maximum Single Application Rate: **DO NOT** exceed the maximum rate listed in the table.
- 2) **DO NOT** apply more than 8.0 fl. oz. of solution by drip/drench application per treatment.
- 3) **DO NOT** apply more than 16.0 fl. oz. of solution by drip/drench application per plant per year.
- 4) Minimum Application Interval: 21 days
- 5) **DO NOT** apply more than 2 applications per crop.
- 6) **DO NOT** apply to the foliage.
- 7) **DO NOT** apply within 1 day of harvest (1-day PHI).
- 8) **DO NOT** use in greenhouse for transplant production.

Crop	Disease	Rate oz/100 gal	Use Directions
Peppers (Greenhouse production only) Pepper, bell Pepper, non-bell (Not for use in California)	Fusarium Root Rot (<i>Fusarium solani</i>)	1.0 oz/100gal water (solution)*	Prepare a drench solution of 1 ounce Cannonball fungicide per 100 gallons water (solution). Apply 5-8 fl. oz. of solution onto the growing medium at the base of each pepper plant.

*1 oz product is equivalent to 0.03 lb ai fludioxonil/100 gal of water

Specific Use Restrictions:

- 1) Maximum Single Application Rate: **DO NOT** exceed the maximum rate listed in the table.
- 2) **DO NOT** apply more than 8 fl. oz. of solution by drip/drench application per treatment.
- 3) **DO NOT** apply more than 1 application per crop.
- 4) **DO NOT** apply to the foliage.
- 5) **DO NOT** apply within 1 day of harvest (1-day PHI).
- 6) **DO NOT** use in greenhouse for transplant production.

Cannonball® and the Syngenta logo are trademarks of a Syngenta Group Company

©2022 Syngenta