

SUPPLEMENTAL LABELING:
HEMP

Carb-O-Nator®



CAN BE USED IN ORGANIC PRODUCTION



ACTIVE INGREDIENT:

Potassium Bicarbonate..... 85.00%

OTHER INGREDIENTS:..... 15.00%

TOTAL 100.00%

Keep Out of Reach of Children
CAUTION

This Supplemental Labeling expires on December 31, 2021 and must not be used or distributed after this date.

MANUFACTURED BY:

Certis USA LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046

EPA Reg. No. 70051-117
EPA Est. No. 70051-CA-001



This is a Specimen Label. It may not reflect the most-recent approved label for use in your state. Always refer to the label on the product packaging for approved use instructions. Please contact your Certis sales representative for more information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in possession of the user at the time of application.

Read the label affixed to the container for **CARB-O-NATOR®** before applying.

Use of **CARB-O-NATOR®** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **CARB-O-NATOR®**

CROP

Hemp

PEST

Gray mold (*Botrytis cinerea*), hemp leaf spot (*Bipolaris* sp.), Powdery mildew (*Leveillula* and *Sphaerotheca* spp.)

APPLICATION RATES

FIELD APPLICATION

Mix at a rate of 2.5 to 5 pounds of **CARB-O-NATOR®** in 100 gallons of water. Apply mixed spray solution to thoroughly wet stems and foliage of plants. Apply a minimum of 20 gallons of mixed spray solution per acre. Spray volume per acre of treatment can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Repeat application at 10 – 14 day intervals or as needed. Increase frequency of application to every 5 – 7 days during periods of weather favorable to fungal growth. **DO NOT EXCEED A MIX RATE OF 5 POUNDS OF CARB-O-NATOR® PER 100 GALLONS OF WATER.** Do not store unused spray solution. Dispose of unused spray solution on site. **Spray solution should be applied within 12 hours of preparation.**

GREENHOUSE APPLICATION

Mix at a rate of 2.5 to 5 pounds of **CARB-O-NATOR®** in 100 gallons of water. Apply mixed spray solution to thoroughly wet stems and foliage of plants. Spray volume per unit of treated area can vary depending upon plant type and stage development. Use minimum of 10–20 gallons of water per 1/10 acre or 4,350 sq. ft. Concentration of spray solution and frequency of application will depend upon disease pressure. Repeat application at 10–14 day-intervals or as needed. Increase frequency of application to every 5–7 days during periods of weather favorable to fungal growth. **DO NOT EXCEED A MIX RATE OF 5 POUNDS OF CARB-O-NATOR® PER 100 GALLONS OF WATER.** Do not store unused spray solution. Dispose of unused solution on site. **Spray solution should be applied within 12 hours of preparation.**

ESL20200617

Ver20200624