

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN APPROVED BY OR SUBMITTED TO EPA FOR APPROVAL.

CARBARYL 4L

2(ee) Recommendation

FOR USE ON TIMOTHY TO CONTROL CEREAL RUST MITE

THIS 2(ee) RECOMMENDATION IS FOR DISTRIBUTION AND USE IN THE STATES OF VIRGINIA AND WEST VIRGINIA.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

This product bulletin contains new or additional directions for use which are recommended by Loveland Products, Inc. under FIFRA Section 2(ee).

DIRECTIONS FOR USE

CROP	PEST	RATE OF APPLICATION
Timothy	Cereal Rust Mite	1 to 1.5 qts/A

Directions for use:

Apply with ground equipment only with adequate water for complete coverage (10 to 50 gallons by ground). Higher water volumes should be used for improved mite control. Apply at approximately 4 weeks after green-up in the Spring. Treatment is recommended in fields with previous history of cereal rust mite and/or when 25% of the plant tillers exhibit curled tips of the new leaf blades within several weeks of green-up. In general, one application at 1 to 1.5 quarts per acre should provide control. If needed, a second application can be made at least 14 days after the first application.

RESTRICTIONS AND PRECAUTIONS

- A maximum of 2 applications per year may be made.
- Do not apply within 14 days of harvest or grazing.
- Do not exceed a total of 1.5 quarts per acre per cutting.
- **BEE CAUTION:** Do not apply this product to target crops or weeds in bloom.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED CARBARYL 4L, EPA REG. NO. 34704-447 LABEL MUST BE FOLLOWED. THE CARBARYL 4L SECTION 3 LABEL AND THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



PO Box 1286
Greeley, CO 80632

3/16/2017

For Agricultural or Commercial Use Only
EPA Reg. No. 34704-447