

Carbol™ Multi-Trace

Carbol™ Multi-Trace is a liquid plant micronutrient solution containing zinc, iron, copper, manganese, boron and molybdenum for use as a foliar feed to maintain or correct micronutrient levels.

0-0-0

Guaranteed Analysis:

Boron (B)	0.03 %	Manganese (Mn)	1.6 %
Copper (Cu)	0.9 %	Molybdenum (Mo)	0.008 %
Iron (Fe)	1.3 %	Zinc (Zn)	3.3 %

Derived from zinc amino acid complex, ferric oxide, copperoxychloride, manganese carbonate, boric acid, sodium molybdate

Net volume = 5.28 gallons (20 l)

Density: (68° F) 10.86 lbs. /gallon (4.93 Kg)

Content • Batch number • Manufacture date



HydroGro LLC.

7201 E. Camelback Rd. # 295

Scottsdale, AZ 85251

Phone: 480-248-9336



Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>



SHAKE WELL • Store at room temperature
Shake well before use • Store at room temperature
Keep out of direct sunlight • Keep container closed when not in use



CAUTION: This product contains Boron (B). The use of this product on any crops other than those recommended may result in serious injury to the crop(s).

CAUTION: This product contains Molybdenum (Mo). The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo). Do not apply to crops which are grazed by ruminant animals.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

CONDITIONS OF SALE: The information contained herein is believed to be accurate and reliable. Buyer/User acknowledges and assumes all liability from use of this material. Follow instructions carefully. Methods of applications, timing, water, soil, water, turf, plant, crop condition, prior control material usage and “acts of God” are beyond the control of seller and manufacturer.

CAUTION: Exercise care when storing, applying and disposing of product. Keep out of reach of children. Follow all proper disposal procedures. Do not ingest.

Carbol™ Multi-trace is compatible with most agricultural remedies. However, do NOT apply **Carbol™ Multi-trace**:

- in combination with copper containing fungicides, phosphates or phosphonates, or within 30 days after application of any copper fungicides.
- in combination with highly alkaline material such as lime sulfur or certain boron products.

Carbol™ Multi-trace should preferably be applied in the early morning or late afternoon. **Do not apply to plants that are undergoing a period of moisture or heat stress.**

- Do not exceed a concentration of 2% (2 gallons/100 gallons water).

For optimum results **Carbol™ Multi-trace** should be used in a program and in combination with other balanced fertilizers and foliar feeds

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Foliarly apply **Carbol™ Multi-trace** according to the following:

CROP	MAX RATE / ACRE	REMARKS
Field & row	3/4-1 quart	Apply at 3-4 leaf stage in 30 gallons water per acre (cereals). For other crops, apply early in growing season in 30- 60 gallons water and repeat before flowering.
Orchard & vineyard	1-2 quart	Start application when shoots are 3-4 inches in length in 100-150 gallons water per acre. Repeat every 10 days for a total of 3-4 applications (vineyards). Start application directly after fruit set, repeat every 10 days for a total of 2-3 applications. Apply in 100-200 gallons water per acre (stone fruit).
Vegetable & fruit	2-3 pints	Start application at fruit set. Repeat every 10 days. Apply in 50 gallons of water per acre.
Other crops	-	Consult your local agronomic professional.

¹ Base your application of **Carbol™ Multi-trace** on petiole or tissue analysis. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.