



carboxy™ Fe

PRECAUTIONARY STATEMENTS

BEFORE USING THIS PRODUCT, READ ALL PRECAUTIONS, DIRECTIONS OF USE.

CAUTION: Avoid contact with formulated product for long periods of time. Do not ingest.

IN CASE OF CONTACT WITH SKIN OR EYES: Rinse with abundant water.

GENERAL INFORMATION

Carboxy Fe™ is an easy to apply product derived from plant extract, **Carboxy Fe™** contains polyhydroxycarboxylic acids. These acids make nutrients more available for root uptake. **Carboxy Fe™** is particularly effective on young developing plants or established perennials beginning active growth in the spring.

STORAGE AND DISPOSAL

Do not transfer into food or drinks containers. The remnant of product should be disposed of in accordance with local requirements.

1. WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. THESE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, THE TYPES OF SOILS, OR THE MANNER OF USE, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER.

Net Weight :

Lot Number:

Date of
Manufacture:

Guaranteed Analysis:

Sulfur (S) 5.1%
Iron (Fe) 9.0%

Derived from Ferrous Sulfate.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

Soil Amending Guaranteed Analysis

Active Ingredients

4.0% Polyhydroxycarboxylic Acids (PHCA)

96% Total Other Ingredients (inert as non-plant food ingredients)

Purpose: Makes nutrients more available for root uptake

Expiration date:

2 years after the date of manufacture.

CAUTION
KEEP OUT OF REACH OF CHILDREN



2. LIMITATION OF LIABILITY

IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. NEITHER MANUFACTURER NOR SELLER SHALL BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, ITS CUSTOMERS, OR USER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT OR RETURN OF THE PRODUCT AT THE SOLE DISCRETION OF MANUFACTURER OR SELLER.

3. If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

© Chemport, Inc. All rights reserved.

DIRECTIONS FOR USE

Apply **Carboxy Fe™** as part of the nutrient solution in fertigation.

CROP	DIRECTIONS FOR USE	
	FERTIGATION	FOLIAR APPLICATION
Vegetables Crops (peppers, tomatoes, bell peppers, eggplant, cucurbits such as watermelon, melon, zucchini, cucumber, crucifer such as cabbage and broccoli, bulb vegetables such as potato, onion and garlic)	Use 1 lb/acre at 10 to 15 days interval during the vegetative growth and 1 to 3 lb/acre at 10 to 15 days interval during fruit set.	Use 0.5 to 1 lb/acre at 15 to 20 days interval.
Berries (blackberry, blueberry, raspberry, strawberry)	Use 1 to 3 lb/acre, 2 to 3 applications during the vegetative growth.	Use 0.5 to 1 lb/acre at 15 to 20 days interval.
Ornamentals	Use 1 to 3 lb/acre, 1 to 2 applications during active vegetative growth depending upon the plant	Use 1 to 2 lb/100 gallon at 15 to 20 days interval.
Temperate Fruit Trees (Apple, Peach and Cherry)	Use 2 to 4 lb/acre, 3 to 5 applications during the vegetative growth and 2 to 3 applications during the fruit set.	Use 1 to 2 lb/acre at 15 to 20 days interval.
Grapes (Table Grape, Raisin Grape and Wine Grape)	Use 2 to 4 lb/acre, 3 to 5 applications during the vegetative growth and 2 to 3 applications during the fruit set.	Use 1 to 2 lb/acre at 15 to 20 days interval.
Nut Crops (Almonds, Walnut and Pistachio)	Use 2 to 4 lb/acre, 3 to 5 applications during the vegetative growth and 2 to 3 applications during the fruit set.	Use 1 to 2 lb/acre at 15 to 20 days interval.
Fruit Trees (Avocado, Mango and Citrus)	Use 2 to 4 lb/acre, 3 to 5 applications during the vegetative growth and 2 to 3 applications during the fruit growth.	Use 0.5 to 2 lb/acre at 15 to 20 days interval.
Hemp	Use 1 lb/acre at 10 to 15 days interval during the vegetative growth.	Use 0.5 to 1 lb/acre every 15 to 20 days interval.

*In case of Fe deficiency, growers are recommended to apply carboxy Fe until symptoms disappears. *In general, spray volume of 100 gallon per acre is recommended to achieve complete crop coverage.

Information about the components of this fertilizer may be obtained by writing to Chemport Inc. and giving the lot number found on this label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

Innovak
GLOBAL

A parent company of: Chemport Inc.
Guaranteed by: Chemport Inc.

1716 Montana Avenue El Paso TX 79902-5705

Phone: (915) 591-49-07