

Cell Power®

Damu™

GUARANTEED ANALYSIS

Boron (B).....9.00%
Molybdenum (Mo)..... 0.10%

Derived from: Boric Acid and Sodium Molybdate

WARNING: This product contains Boron(B) and may be harmful to certain crops. Use only according to manufacturer's directions. **WARNING:** The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

F2737

KEEP OUT OF REACH OF CHILDREN



WARNING

Hazard Statements: Causes skin irritation. Causes serious eye irritation.

Precautionary Statements: Wash hands thoroughly after handling. Wear protective gloves. Wear eye, clothing and face protection.

FIRST AID: **If on skin:** Wash with plenty of soap and water. If skin irritation occurs, get medical advice or attention. Take off contaminated clothing and wash it before reuse. **If in eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice or attention.

FOR AGRICULTURAL USE ONLY

V:081221

GUARANTEED BY: OMEX® AGRIFLUIDS, INC. • PO BOX 815 • SELMA, CA 93662



DENSITY: 10.85 LBS/GALLON @ 68° F

LIQUID CONTENT 2.5L US GALLONS (9.5 L) NET WEIGHT: 27.13 LBS (12.3 kg)
LIQUID CONTENT 265L US GALLONS (1,003L) NET WEIGHT: 2875 LBS (1304 kg)



GENERAL INFORMATION

Cell PowerDamu can be used in combination with water soluble plant nutritional products which are formulated for drip and foliar and drip application. This product may be applied alone or in tank mixes

containing. **Cell PowerDamu** is believed to be compatible with other nutritional products, but when in doubt a jar test should be used to ensure compatibility. When applying **Cell PowerDamu** as foliar, apply with sufficient water to provide good coverage on all crops.

ONCE PRODUCT IS DILUTED, USE WITHIN 24 HOURS

DIRECTIONS FOR USE

All Permanent Crops including all Stone Fruit, Pome Fruits and Tree Nuts, such as Almond, and *Pistachio: One application pre-bloom at 16 oz. per acre may be applied. Apply post-harvest applications of up to 32 oz/acre prior to true dormancy. Apply one treatment foliar or drip at 32 oz/acre two weeks prior to harvest.

STONE FRUIT (Cherry, Apricot, Plum, Nectarine): Apply 16 oz/acre at flowering. You may make a second application at post-harvest, but prior to leaf full at a rate of 32 oz/acre. **Strawberry:** Apply foliar or drip 8-16 oz/acre every 2-3 weeks during flowering and fruiting period.

Caneberries, (including, but not limited to Blackberries, Blueberries, Cranberries and Raspberries): Apply as a foliar or irrigation, 16 oz/acre at fruit set and then a second application 1-2 weeks prior to harvest at 16 oz/acre.

All Fruit crops including, but not limited to Melons, Peppers, Eggplant and Tomatoes: Apply as foliar or irrigation, 16 oz/acre at fruit set and then a second application 1-2 weeks prior to harvest at 16 oz/acre.

All Leafy Vegetable and Cole crops (including, but not limited to all the root vegetables Asparagus, Tomatoes): Apply 16 oz/acre through irrigation 4-6 leaf stage, repeat every 4-6 weeks. **CUCURBIT VEGETABLES such as Cantaloupe, Watermelon, Citron Melon, Cucumbers and Pickles:** Apply 16-32 oz/acre irrigation during flowering periods. Apply 64 oz/acre foliar or drip 7-14 days prior to harvest. **Citrus:** Apply 16 oz/per acre after each leaf flush. May be applied foliar or drip.

Cole Crops, such as Broccoli, Brussels Sprouts, Cauliflowers: Apply 16 oz/acre via irrigation 4-6 leaf stage, repeat every 4-6 weeks.

Cereal Grains such as Hops: Apply 8 oz/acre foliar between flowering and milky ripe stage development.

Cotton: Full season, Irrigated Cotton: Apply 16 oz/acre foliar combine with PIX applications at 16 oz/acre monthly applications from squaring and use less PIX. Apply 16 oz/acre foliar during main flowering.

All Fresh and Dried Beans (including but not limited to Soybeans, Lima and Snap Beans): At flowering apply foliar 8oz/acre. Make a second application at 8oz/acre at pod fill with sufficient water for good coverage.

Tropical Tree Fruits such as Avocado, mango and guava: Apply 8-16 oz/acre foliar monthly applications.

Grapes (Wine) and Grapes, such as Dried, Raisin, Seedless, and Seeded Table Grapes: Apply 16 oz/acre. For seedless and seeded grapes apply after thinning and bloom set has passed. Apply second application 20 days after fruit set 1-2 weeks ahead of harvest. Apply all applications through the irrigation system. **Root Vegetables:** Apply 16-32 oz/acre as a foliar application. Apply 64 oz/acre foliar 7-14 days prior to harvest.

Sugar beets and Potatoes: Apply 16-32 oz/acre for irrigation or foliar applications when growing conditions are generating rapid shoot growth to address nutrient deficiency. Apply 64 oz/acre as a foliar 7-14 days prior harvest.

STORAGE and DISPOSAL

STORAGE: Store in a cool, dry, well-ventilated place out of direct sunlight. Store in original tightly closed container away from children, animals, foods, feeds, and seeds. Store at ambient temperatures.

Product and Container Disposal: Do not contaminate water, food, or feed by storage or disposal. Empty containers or liners may retain some product residues. Triple rinse (or equivalent), during mixing and loading. Dispose of contents in accordance with local/regional/national/international regulations.

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale-Warranty and Limitations of Liability and Remedies before using this product. The directions should be followed carefully together with good application practice. Failure to do so may cause injury to crops, animals, man, or the environment. Extraordinary or unusual weather conditions may affect product performance and/or crop injury which are beyond the control of Omex Agrifluids, Inc. (the "Company") or seller. The Company warrants that this product conforms to the chemical description on the label and is reasonably for the purpose referred to in the Directions For Use. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Company's election, ONE of the following: 1. Refund of the purchase price paid by buyer or user for product bought. 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The company/seller offer this product and the buyer/user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability, and remedies.

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals/html>.

