

Cell Power®

SizeN® Multi Mix

15-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
11.0% Urea Nitrogen	
4.00% Nitrate Nitrogen	
Calcium (Ca)	5.40%
Magnesium (Mg)	0.50%
0.5% Water Soluble Magnesium (Mg)	
Copper (Cu)	0.05%
0.05% Water Soluble Copper	
Iron (Fe)	0.10%
0.10% Water Soluble Iron	
Manganese (Mn)	0.50%
0.50% Water Soluble Manganese	
Zinc (Zn)	0.30%
0.3% Water Soluble Zinc (Zn)	

Derived From: Urea, Calcium Nitrate, Magnesium Chloride Hexahydrate, Copper Chloride Dihydrate, Manganese Chloride Tetrahydrate, Iron Chloride, Zinc Chloride Anhydrous

F2737

KEEP OUT OF REACH OF CHILDREN

ANGER



Hazard Statements: Harmful if swallowed. Causes serious eye damage.

Precautionary Statements: Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Wear eye and face protection. Wear protective gloves.

FIRST AID: If **swallowed**: Call a poison control center or doctor if you feel unwell. Rinse mouth. If **in eyes**: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a poison control center or doctor.

FOR AGRICULTURAL OR COMMERCIAL

V:0081221

GUARANTEED BY: OMEX® AGRIFLUIDS, INC. • PO BOX 815 • SELMA, CA 93662



DENSITY: 11.00 LBS./GALLON @ 68° F

LIQUID CONTENT 2.5 GALLONS (9.46 L) NET WEIGHT: 27.50 LBS. (12.47 kg)

LIQUID CONTENT 264 GALLONS (1,003 L) NET WEIGHT 2,904 LBS (1317 kg)

GENERAL INFORMATION

Cell Power SizeNMulti Mix is a unique blend of nitrogen, calcium and micronutrients. **Cell Power SizeNMulti Mix** can be used in combination with water soluble plant nutritional products which are formulated for drip and foliar and drip application. This product may be applied alone or in tank mixes. **Cell Power SizeNMulti Mix** is believed to be compatible with other nutritional products, but when in doubt a jar test should be used to ensure compatibility. When applying **Cell Power SizeNMulti Mix** as foliar, apply with sufficient water to provide good coverage on all crops.

ONCE PRODUCT IS DILUTED, USE WITHIN 24 HOURS

DIRECTIONS FOR USE

Dissolve the desired amount of **Cell Power SizeNMulti Mix** in water before adding to the spray tank. With proper agitation, **Cell Power SizeNMulti Mix** can be added into the spray tank with agitation running.

SUGGESTED USE RATES

CROP - RATE/ACRE - NUMBER & TIMING OF APPLICATIONS ALL

Permanent Crops: Apply 1-2 quarts/acre near bloom. Continue 1-2 quarts/acre applications every two weeks during the fruit sizing period. One application pre-bloom at 1-2 quarts/acre per acre may be applied. Apply post-harvest applications of up to 1-4 quarts/acre prior to true dormancy on all permanent crops. May be applied through irrigation water.

All Strawberries, Caneberries & Blueberries: Begin applying as early as bloom at 1-2 quarts/acre. Continue the applications every two weeks during the flowering/fruit setting cycles of the crop. For Blueberries which are more determinate, begin applications at fruit set and go every two weeks during the fruit sizing period. May be applied through irrigation water.

All Field Crops and Hops: Begin application of 1-2 quarts/acre prior to bloom and continue for two weeks during the active growing prior of the crop. May be applied through irrigation water.

All Vegetable Crops: Begin applications right after thinning/planting and make 2-4 total applications at 1-2 quarts/acre every two weeks during the most active period of the crop's growth.

Make all applications with sufficient water for good coverage. Irrigation water application: This product may be applied at the labeled rates through the following irrigation systems: Furrow, drip/micro, center-pivot, solid-set, side-roll, border check. A jar test with source water and/or other fertilizers should be initiated to ensure compatibility.

Apply with sufficient water for good coverage. For higher volume sprays, rate should be adjusted. Caution: Store **Cell Power SizeNMulti Mix** in temperatures between 41° F to 104° F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product. Use caution when applying to fruiting crops in combination with pesticides and/or surfactants. Use minimal effective rates of stickers during ripening. Do not use high analysis organo silicones or high analysis non-ionics during ripening. Avoid application to fruit at elevated temperatures (>95° F). Avoid applications to crops under environmental stress or pest pressure. Maximum effectiveness will be obtained when applied early in the morning or after dusk.

STORAGE and DISPOSAL

Storage: Store in a cool, dry, well-ventilated place out of direct sunlight. Store in original tightly closed container away from children, animals, foods, feeds, and seeds. Store at ambient temperatures.

Product and Container Disposal: Do not contaminate water, food, or feed by storage or disposal. Empty containers or liners may retain some product residues. Triple rinse (or equivalent), during mixing and loading. Dispose of contents in accordance with local/regional/national/international regulations.

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale-Warranty and Limitations of Liability and Remedies before using this product. The directions should be followed carefully together with good application practice. Failure to do so may cause injury to crops, animals, man, or the environment. Extraordinary or unusual weather conditions may affect product performance and/or crop injury which are beyond the control of OMEX Agrifluids, Inc. (the "Company") or seller. The Company warrants that this product conforms to the chemical description on the label and is reasonably for the purpose referred to in the Directions For Use. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Company's election, ONE of the following: 1. Refund of the purchase price paid by buyer or user for product bought. 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The company/seller offer this product and the buyer/user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability, and remedies.

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

