

# CELL POWER<sup>®</sup>

## Tru Boost<sup>®</sup> 5-0-0

### GUARANTEED ANALYSIS

|                               |       |
|-------------------------------|-------|
| Total Nitrogen (N) .....      | 5.00% |
| 5.00% Urea Nitrogen           |       |
| Magnesium (Mg) .....          | 2.40% |
| 2.40% Water Soluble Magnesium |       |
| Sulfur (S) .....              | 3.20% |
| 3.20% Combined Sulfur (S)     |       |
| Boron (B) .....               | 2.00% |
| Molybdenum (Mo) .....         | 0.02% |

**Derived from:** Urea, Magnesium Sulfate, Boric Acid, and Sodium Molybdate

**WARNING:** This product contains boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions.

**KEEP OUT OF REACH OF CHILDREN**



## WARNING

**Hazard Statement:** Suspected of damaging fertility or the unborn child.

**Precautionary Statements:** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing and eye/face protection.

**FIRST AID:** If exposed or concerned, get medical advice or attention.



**FOR AGRICULTURAL USE ONLY**

V:033021

GUARANTEED BY: OMEX<sup>®</sup> AGRIFLUIDS, INC. • PO BOX 815 • SELMA, CA 93662

**DENSITY: 10.50 LBS/GALLON @ 68° F**

**LIQUID MEASURE 2.5 US GALLONS (9.46 L) NET WEIGHT: 26.25 LBS(11.90 kg)**

**LIQUID MEASURE 265 US GALLONS (1003 L) NET WEIGHT: 2783 LBS(1262 kg)**

**CAUTION:** This fertilizer is to be used only on soil which responds to molybdenum. Crops high in molybdenum are toxic to grazing animals (ruminants).

**CAUTION:** This product contains Boron (B) and may be harmful to certain crops. Use only according to manufacturer's directions.

## DIRECTIONS FOR USE

### VEGETABLE CROPS

**Fruit-bearing vegetables:** For fruit-bearing vegetable crops not specifically listed on label, apply **Tru Boost** at pre-bloom, bloom and every 1 to 2 weeks thereafter at a rate of 3 pints to 2 quarts per acre.

**CUCURBITS, PEPPERS, TOMATOES, AND OTHER FRUITING VEGETABLES:** 3 pints per acre per application. 1st application at first flower. 2nd application 1 to 2 weeks later. 3rd application 1 to 2 weeks later.

**POTATOES** 3 pints per acre per application. 1st application at 100% sprouting. 2nd application 15 days after first treatment.

**EDIBLE BEANS, PEAS, SNAPBEANS, & OTHER LEGUMES:** 1 quart per acre per application: 1st application at early bloom with an optional 2nd application at early pod.

**SWEET POTATOES:** 3 pints per acre per application. 1st application at Tuber initiation. 2nd application 2 to 3 weeks after first treatment.

**SUGARBEET:** 3 pints per acre per application. 1st application at 8 to 10 leaf stage followed by a 2nd application 2 to 3 weeks later.

### ROW & NUT CROPS

**COTTON:** 1 quart per acre per application. Apply at early to mid bloom.

**SOYBEAN:** 2 to 3 pints per acre per application. Apply at R1 to R3.

**PEANUTS:** 1 quart per acre per application. 1st application at pegging. 2nd application 10 to 14 days after.

**ALMONDS & PISTACHIOS:** 2 quarts per acre per application. 1st application at pre-bloom to early bloom. 2nd application at full bloom to petal fall, or 1 to 3 quarts per acre application at full bloom.

**ENGLISH WALNUTS AND PECANS:** 2 quarts per acre per application. 1st application at early pistillate bloom. 2nd application at full pistillate bloom, or 1 to 3 quarts per acre application at full pistillate bloom.

### FRUIT CROPS

Apply **Tru Boost** to most fruit crops at pre-bloom, full bloom and soon after fruit set at a rate of 1 quart, 3 pints or 2 quarts per acre depending on the fruit crop.

**APPLES & PEARS:** 3 pints per acre per application. 1st application at pre-bloom. 2nd application at early king bloom (for apples); bloom (for pears). 3rd application at post bloom (petal fall-fruit set).

**STONEFRUIT (Apricots, Nectarines, Peaches, Plums):** 3 pints per acre per application. 1st application at early bloom. 2nd application at petal fall. 3rd application 10 to 20 days later (dependent upon variety).

**GRAPES (Raisin, Table and Wine):** 1 quart per acre per application. 1st application at 12 to 18 inch shoot growth. 2nd application at bloom stage. 3rd application after fruit set.

**Concord, Niagara and other American cultivars:** 2 quarts per acre per application. 1st application at early bloom. 2nd application at full bloom.

**POMEGRANATE:** 2 quarts per acre per application. 1st application at early bloom. 2nd application at petal fall.

**CHERRIES:** 2 quarts per acre per application. 1st application at first white flower. 2nd application at petal fall. 3rd application at shuck fall.

**OLIVES:** 3 pints per acre per application. 1st application at bud differentiation. 2nd application at early to full bloom. 3rd application at first growth stage of fruit.

**STRAWBERRIES:** 1 quart per acre per application. 1st application within 14 days of transplant. 2nd application at first bloom. 3rd application 2 to 3 weeks later. Subsequent applications every 2 to 3 weeks thereafter.

**BLUEBERRY, CANEBERRY AND CRANBERRY:** 1 quart per acre per application. 1st application at swollen bud. 2nd application at first bloom. 3rd application at full bloom. 4th application at petal fall.

**AVOCADOS:** 2 quarts per acre per application. 1st application at pre-bloom to early bloom. 2nd application at full bloom to petal fall.

**CITRUS:** 2 quarts per acre per application. 1st application at pre-bloom to early bloom. 2nd application at full bloom to 2/3 petal fall. 3rd application at fall spray (late August for early varieties to late October for late varieties).



## OIL SEED CROPS

Apply **Tru Boost** to most oil seed crops at budding to early bloom at a rate of 1.5 pints to 2 pints per acre.

**CANOLA:** 1.5 pints per acre per application. One application at beginning of stem elongation.

**CAMELINA:** 2 pints per acre per application. One application at budding to early flower.

**FLAX:** 2 pints per acre per application. One application before onset of flowering.

**MUSTARD:** 2 pints per acre per application. One application at budding to early flower

**SUNFLOWER:** 2.5 pints per acre applied at 3 pairs of true leaves to flower bud appearance.

## SOIL APPLICATION

Apply 2 to 3 pints per acre anytime during the growing season. Repeat as needed. May be applied via fertigation or with all conventional spray equipment including side dress application. At least one post-harvest application is recommended for all permanent crops.

## STORAGE and DISPOSAL

**STORAGE:** Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal. **DISPOSAL:** Do not reuse container. Plastic: Triple rinse (or equivalent), adding rinse water to spray tank, then offer for recycling at an ACRC site (go to <http://www.acrecycle.org/> for locations) or by reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of OMEX® AGRIFLUIDS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of OMEX® AGRIFLUIDS, INC. and the seller. The buyer or use of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, OMEX® AGRIFLUIDS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS", AND OMEX® AGRIFLUIDS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT OMEX® AGRIFLUIDS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: OMEX® AGRIFLUIDS, INC., ATTENTION LAW DEPARTMENT, PO BOX 815, SELMA, CA 93662.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING: AT THE ELECTION OF OMEX® AGRIFLUIDS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OMEX® AGRIFLUIDS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>