

SUPPLEMENTAL LABELING

BUPROFEZIN	GROUP	16	INSECTICIDE
------------	-------	----	-------------



EPA Reg. No. 71711-21

For Use on Bushberry (Crop Group 13-07B) – Not for Use in California

This supplemental label expires 4/1/2025 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for Centaur® Insect Growth Regulator before applying. Use of Centaur Insect Growth Regulator according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Centaur Insect Growth Regulator.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART

Bushberry (Crop Subgroup 13-07B) – Not for Use in California		
aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry, native currant; salal; sea buckthorn; cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Scales Leafhoppers	34.5 oz (1.5 lb ai)	<ul style="list-style-type: none">• Apply by ground application using a minimum of 30 gallons of water per acre. <p>USE RESTRICTIONS</p> <ul style="list-style-type: none">• Do not apply more than 34.5 oz of product per acre per growing season.• Do not make more than 1 application per growing season.• Preharvest Interval (PHI): 60 days <p>RECOMMENDATIONS</p> <ul style="list-style-type: none">• Thorough spray coverage is essential.• San Jose scale: Application can be made at dormant or delayed dormant when scale is in the black cap stage or in-season at early crawler emergence.• Other scales: Apply at early crawler emergence or when scales are at a susceptible early nymphal stage.• Leafhoppers: Apply when majority of the population is at the early nymphal stage.

©2023 Nichino America, Inc. Centaur is a registered trademark of Nichino America, Inc.



Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700