
SUPPLEMENTAL LABEL



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

SUPPLEMENTAL LABEL FOR CENTRIC[®] 40WG INSECTICIDE

Supplemental Directions for Use on Cotton

EPA Reg. No. 100-1147

Active Ingredient:

Thiamethoxam¹40.0%

Other Ingredients:60.0%

Total:100.0%

¹CAS No. 153719-23-4

Centric 40WG is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

This label expires July 1, 2012

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Centric 40WG Insecticide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling supersedes the Directions for Use on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SCP 1147S-L1 1106

Centric[®] and the Syngenta logo are trademarks of a Syngenta Group Company

CROP USE DIRECTIONS

Recommended Applications

Crop	Pest	Rate Per Acre Per Application	Remarks
Cotton	Aphids	1.25-2.0 oz./A	Aphids: For best results, the low rate should not be used after first bloom or on rapidly increasing populations.
	Cotton fleahopper Clouded plant bug Tarnished plant bug (<i>Lygus lineolaris</i>)	1.25-2.5 oz./A	Cotton fleahopper, Clouded plant bug and Tarnished plant bug: Control may require the use of two applications made at a 5 – 10 day interval.
	Brown stink bug (suppression only) Green stink bug Southern green stink bug Thrips Whiteflies	2.0-2.5 oz./A	Stink bugs, Thrips, and Whiteflies: Control may require the use of two applications made at a 5 – 10 day interval.

Remarks:

- **Application Timing:** Apply before pests reach damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Apply higher rates for heavy infestations. Do not apply this product within 45 days of planting if cottonseeds were treated with a neonicotinoid product.
- **Pre-Harvest Interval (PHI):** 21 Days.
- **Maximum Centric Allowed per Growing Season:** Do not exceed a total of 5.0 oz. of Centric 40WG (0.125 lbs. a.i.) /A during the growing season.
- **Minimum Interval Between Applications:** Allow at least 5 days between applications.
- **Water Volume:** Use sufficient water volume to ensure thorough coverage of foliage. Do not use less than 10 GPA for ground applications and 3 GPA for aerial applications.