

SPECIMEN LABEL

CEREALSTAR™ SELECT ST

Mefenoxam	GROUP	4	FUNGICIDE
Ipconazole	GROUP	3	FUNGICIDE
Thiamethoxam	GROUP	4A	INSECTICIDE

ACTIVE INGREDIENTS:

Thiamethoxam	11.20%
Mefenoxam	1.45%
Ipconazole	0.43%
OTHER INGREDIENTS:	86.92%
TOTAL:	100.00%

Contains 1.00 pound Thiamethoxam per gallon (120.0 grams per liter),
0.13 pound Mefenoxam per gallon (15.6 grams per liter), 0.04 pound
Ipconazole per gallon (4.8 grams per liter)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use inside booklet.

Seed Treatment

*A seed treatment for protection against listed insect
pests and seed/soilborne diseases of Barley, Wheat,
Oats, Rye, Triticale.*

Manufactured By:
ALBAUGH, LLC
1525 NE 36th Street, Ankeny IA 50021



**COMBINATION
SEED TREATMENT**

EPA Reg. No. 45002-41
AD090622

 **ALBAUGH®**
your alternative

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling poison control center (888-347-6732) or doctor, or going for treatment. For 24- Hour Emergency Assistance, chemical spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils

Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and aquatic invertebrates. For terrestrial use, DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment wash water or rinsate. Treated seed exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading and planting.

Pollinator Advisory:

Thiamethoxam is highly toxic to bees and effects may be possible as a result of exposure to translocated residues in blooming crops. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Groundwater Advisory:

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory:

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of mefenoxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is for use in commercial seed treatment facilities or as an end-use seed treatment on agricultural establishments at, or immediately before planting subject to the Restrictions listed below and as specified for each crop in the Crop Specific Use Instructions on this label.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 48 hours**. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

Coveralls, Chemical-resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils, Shoes plus socks, Protective eyewear.

PRODUCT INFORMATION

This product is a systemic and contact broad-spectrum fungicide plus systemic insecticide, ready to use seed treatment containing mefenoxam / ipconazole fungicides and thiamethoxam insecticide. Ipconazole and mefenoxam protect seed against a wide variety of soil borne and seed borne diseases as listed in the seed treatment table below including *Fusarium*, *Cochliobolus sativus*, *Pythium*, smuts, bunt and leaf stripe. It is also effective against seed rot fungi such as *Aspergillus* and *Penicillium*. Thiamethoxam provides early season protection against listed insect pests such as aphids (including bird cherry-oat, English grain, greenbug, and Russian wheat aphid). This product provides early season suppression of Hessian fly damage and wireworm activity and provides protection from damage caused by white grubs (including northern masked chafer, European chafer, and Japanese beetle). When a rate range is given, use higher rates of this product when conditions favor higher disease pressure. Seed should be sound and well cured before treatment.

USE RESTRICTIONS

- **DO NOT** use as a planter (hopper) box treatment.
- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **DO NOT** make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with CerealStar™ Select ST.
- The maximum application rate for thiamethoxam (including seed treatments, foliar applications, soil applications) is 0.266 lbs. AI per acre per year. **DO NOT** apply more than 0.266 lbs. thiamethoxam ai per acre per year.
- The maximum application rate for mefenoxam- and metalaxyl- (including seed treatments, foliar applications, soil applications) is 12.3 lbs. AI per acre per year. **DO NOT** apply more than 12.3 lbs. mefenoxam ai per acre per year.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to wildlife. Cover or incorporate spilled treated seed. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- **Rotational Crop Intervals:**
 - Alfalfa, Brassica (cole) leafy vegetables, buckwheat, corn, pearl millet, proso millet, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice, cotton, cucurbit vegetables, dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables, mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, peanuts, rapeseed seed, and safflower seed), root vegetables, strawberry, sunflowers, tobacco, and tuberous and corm vegetables may be planted 30 days from the date the CerealStar™ Select ST treated seed was planted.
 - For any other crop, the minimum plant back interval is 120 days from the date the CerealStar™ Select ST treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- If this product is undyed, the purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as a food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye added to treated seed must be cleared for use under 40 CFR, Part 180.1001.

RESISTANCE MANAGEMENT

Thiamethoxam	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

For insecticide resistance management, this product contains a Group 4A insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists.

To delay insecticide resistance, take the following steps:

- Rotate the use of this product or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross- resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

For additional information on Insecticide Resistance Management:

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org>

Mefenoxam	GROUP	4	FUNGICIDE
Ipconazole	GROUP	3	FUNGICIDE

For fungicide resistance management, this product contains ipconazole, a Group 3 fungicide; and mefenoxam, a Group 4 fungicide; Ipconazole belongs to the triazole class of chemistry and is a demethylation inhibitor of sterol biosynthesis (DMI) which disrupts membrane synthesis of the fungal cell. Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis.

Fungal populations may contain individuals naturally resistant to Group 3, or 4 fungicides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as alternation with fungicides with a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Albaugh encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact your local extension specialist or certified crop advisor
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>

SEED TREATMENT

This product is a combination of Thiamethoxam insecticide and a mixture of the fungicides Mefenoxam and Ipconazole. When used on listed crops, the fungicide mixture in this product provides consistent control when disease pressure is high. This product aids in the control or suppression of the following insects pests and soilborne diseases of seed for listed crops.

CROP	RATE*	DISEASE CONTROL	INSECT CONTROL
Wheat ¹	5.0 fl.oz. / 100 lbs. seed.	General seed rots (including those caused by <i>Penicillium</i> and <i>Aspergillus</i>), Early season disease control of <i>Pythium</i> damping-off, Stinking smut, Flag smut, Loose smut, Early season <i>Septoria</i> disease complex, Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and early season suppression of wheat leaf rust.	Early season protection against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), White grubs (including European and northern masked chafer, and Japanese beetle), Hessian fly (suppression only), and Wireworms.
Barley ²	5.0 fl.oz. / 100 lbs. seed.	General seed rots (including those caused by <i>Penicillium</i> and <i>Aspergillus</i>), Early season disease control of <i>Pythium</i> damping-off and Covered smut, True Loose smut, Barley stripe (suppression), Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and Early season suppression of barley leaf rust.	
Oats ³	5.0 fl.oz. / 100 lbs. seed.	General seed rots (including those caused by <i>Penicillium</i> and <i>Aspergillus</i>), Early season disease control of <i>Pythium</i> damping-off, Stinking smut, Flag smut, Loose smut, Early season <i>Septoria</i> disease complex, Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and early season suppression of wheat leaf rust.	
Rye ⁴ Triticale ⁵	5.0 fl.oz. / 100 lbs. seed.	General seed rots (including those caused by <i>Penicillium</i> and <i>Aspergillus</i>), Early season disease control of <i>Pythium</i> damping-off, Stinking smut, Flag smut, Loose smut, Early season <i>Septoria</i> disease complex, Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and early season suppression of wheat leaf rust.	

SEED TREATMENT *(continued)*

APPLICATION DIRECTIONS: This product is formulated both for commercial and for on-farm application. Apply this product only with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease and insect protection and optimum performance. For best activity of this product, seed must be sound and well-cured before treatment. Ensure thorough coverage.

*ACTIVE INGREDIENT USE RATE CONVERSIONS

Active Ingredient	Grams AI / 100 KG Seed	MG AI / Seed				
		Wheat	Barley	Oats	Rye	Triticale
Thiamethoxam	8.04	0.0118	0.0127	0.0148 – 0.0104	0.0098	0.0154 – 0.0135
Mefenoxam	1.05	0.0015	0.0016	0.0019 – 0.0014	0.0013	0.0020 – 0.0018
Ipconazole	0.40	0.0005	0.0006	0.0007 – 0.0005	0.0005	0.0008 – 0.0007

¹ Based on 15,000 seeds / lb

² Based on 14,000 seeds / lb

³ Based on 12,000 – 17,000 seeds / lb

⁴ Based on 18,000 seeds / lb

⁵ Based on 11,500 – 13,000 seeds / lb

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with CerealStar™ Select ST, EPA Reg. No. 45002-UR (thiamethoxam, mefenoxam, ipconazole).
- **DO NOT** use for feed, food, or oil purpose.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH CEREALSTAR™ SELECT ST (thiamethoxam, mefenoxam, ipconazole):

- **Pollinator Precautions:** Thiamethoxam is highly toxic to bees and effects may be possible as a result of exposure to translocated residues in blooming crops. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- **Groundwater Advisory:** Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow. Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- **Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of mefenoxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.
- Store away from food and feedstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to wildlife. Cover or collect spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

- **Rotational Crop Intervals:** Alfalfa, Brassica (cole) leafy vegetables, buckwheat, corn, pearl millet, proso millet, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice, cotton, cucurbit vegetables, dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables, mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, peanuts, rapeseed seed, and safflower seed), root vegetables, strawberry, sunflowers, tobacco, and tuberous and corm vegetables may be planted 30 days from the date the CerealStar™ Select ST treated seed was planted. For any other crop, the minimum plant back interval is 120 days from the date the CerealStar™ Select ST treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for thiamethoxam is 0.266 lbs.
- This seed has been treated with __X__ lbs. of thiamethoxam, __X__ lbs. of mefenoxam, and __X__ lbs. of ipconazole.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

CONTAINER HANDLING:

[less than or equal to 5 gallons]

Non-refillable HDPE container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

[greater than 5 gallons]

Non-refillable HDPE container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

[refillable containers]

Refillable HDPE container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

CONDITIONS OF SALE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

ALBAUGH, LLC and the Seller offer this product subject to the following Conditions of Sale, Warranty Disclaimer and Limitation of Liability, and by purchasing and using this product, Buyer and Users accept such Conditions of Sale, Warranty Disclaimer and Limitation of Liability. No employee or agent of ALBAUGH, LLC or the Seller is authorized to modify the terms of these Conditions of Sale, Warranty Disclaimer and Limitation of Liability unless in writing and signed by a duly authorized representative of ALBAUGH, LLC.

The **DIRECTIONS FOR USE** of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC or the Seller. All such risks shall be assumed by the Buyer. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**

TO THE FULLEST EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR ANY OTHER EXPRESS OR IMPLIED WARRANTY THAT EXTENDS BEYOND THE STATEMENTS MADE ON THIS LABEL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S AND SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH, LLC OR THE SELLER DISCLAIMS ANY LIABILITY FOR INCIDENTAL, CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**