

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1530A-S1 0915**

<b>GROUP</b>	<b>3</b>	<b>12</b>	<b>FUNGICIDES</b>
--------------	----------	-----------	-------------------

**Chairman™**

Fungicide

**This supplemental label expires on 09/22/2018 and must not be used or distributed after this date.**

Active Ingredients:

Fludioxonil:* .....	21.7%
Propiconazole:** .....	9.3%
<hr/>	
Other Ingredients:	69.0%
<hr/>	
Total:	100.0%

\*CAS No. 131341-86-1

\*\*CAS No. 60207-90-1

Chairman is a suspoemulsion (SE) formulation.

Chairman contains 2.03 lb ai per gallon of fludioxonil and 0.86 lb ai per gallon of propiconazole.

**KEEP OUT OF REACH OF CHILDREN.**

## **CAUTION**

**EPA Reg. No. 100-1530**

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Chairman as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**



## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

---

### CROP USE DIRECTIONS

---

**Stone Fruit:** Apricot (*Prunus armeniaca*); Nectarine (*Prunus persica*); Peach (*Prunus persica*); Plum (*Prunus domestica*, *Prunus* spp.); Plum, Chickasaw (*Prunus angustifolia*); Plum, Damson (*Prunus domestica* spp. *insititia*); Plum, Japanese (*Prunus salicina*); Plumcot (*Prunus armeniaca* × *P. domestica*); Prune (fresh) (*Prunus domestica*, *Prunus* spp.); as well as other cultivars and hybrids of these

Use Chairman as a post-harvest dip, drench or spray for the control of post-harvest diseases caused by: *Monilinia* spp., *Botrytis cinerea*, *Rhizopus stolonifer*, *Gilbertella persicaria*, and *Geotrichum* spp.

Application Method	Disease	Rate (fl oz)	Remarks
In-Line Dip/Drench Application <b>For Nectarines ONLY</b>	Brown rot Gray mold Rhizopus rot Gilbertella rot Sour rot	16 fl oz/100 gal	<ul style="list-style-type: none"> <li>Mix 16 fl oz of Chairman in 100 gallons of water, wax/emulsion, or aqueous dilution of wax/oil emulsion.</li> <li>Dip for approximately 30 seconds and allow fruit to drain.</li> </ul>
In-Line Aqueous or Fruit Coating Spray Application	Brown rot Gray mold Rhizopus rot Gilbertella rot Sour rot	16 fl oz/200,000 lb of fruit	<ul style="list-style-type: none"> <li>Ensure proper coverage of the fruit.</li> <li>Mix 16 fl oz of Chairman in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated.</li> <li>Use T-Jet, CDA, or similar application system.</li> <li>For maximum efficacy, use low volume concentrate application systems for treatment of plums.</li> </ul>

**Restrictions:**

**Do not make more than one post-harvest application to the fruit.**

- Ensure the Chairman solution remains in suspension by using agitation.

**Cherries:** Cherry, sweet (*Prunus avium*); Cherry, tart (*Prunus cerasus*); as well as other cultivars and hybrids of these

Application Method	Disease	Rate (fl oz)	Remarks
In-Line Dip/Drench Application	Brown rot Gray mold Rhizopus rot Gilbertella rot Sour rot	16 fl oz/100 gal	<ul style="list-style-type: none"> <li>Mix 16 fl oz of Chairman in 100 gallons of an appropriate water, wax/emulsion, or aqueous dilution of a wax/oil emulsion.</li> <li>Dip for approximately 30 seconds and allow fruit to drain.</li> </ul>
In-Line Aqueous or Flooder Application  High-Volume (Dilute-Spray) Application	Brown rot Gray mold Rhizopus rot Gilbertella rot Sour rot	16 fl oz/50,000 lb of fruit	<ul style="list-style-type: none"> <li>Ensure proper coverage of the fruit.</li> <li>Mix 16 fl oz of Chairman in an appropriate water, wax/emulsion, or aqueous dilution of a wax/oil emulsion.</li> <li>Use flooders, T-jet, or similar application system.</li> </ul>

**Restrictions:**

**Do not make more than one post-harvest application to the fruit.**

- Ensure the Chairman solution remains in suspension by using agitation.

## Tomato

Use Chairman as a post-harvest dip, drench or high volume spray for the control of certain post-harvest rots caused by *Alternaria alternata*, *Botrytis cinerea*, *Rhizopus stolonifer* and *Geotrichum candidum* on tomatoes.

<b>Tomato – Large and Medium Fruiting Varieties</b>			
<b>Application Method</b>	<b>Disease</b>	<b>Rate (fl oz)</b>	<b>Remarks</b>
In-Line Dip/Drench Application	Black mold Gray mold Rhizopus rot Sour rot	16-32 fl oz/100 gal	Mix 16-32 fl oz of Chairman in 100 gallons of water or an appropriate water, wax/emulsion, or aqueous dilution of wax/oil emulsion.  Dip for approximately 30 seconds and allow fruit to drain.
High-Volume (Dilute Spray) Application	Black mold Gray mold Rhizopus rot Sour rot	16-32 fl oz/50,000 lb of fruit	Ensure proper coverage of the fruit.  Mix 16-32 fl oz of Chairman in an appropriate water, wax/emulsion, or aqueous dilution of a wax/oil emulsion for the crop treated.  Use T-jet or similar application system.
<b>Tomato – Small Fruiting Varieties (&lt;1 inch in diameter)</b>			
<b>Application Method</b>	<b>Disease</b>	<b>Rate (fl oz)</b>	<b>Remarks</b>
In-Line Dip/Drench Application	Black mold Gray mold Rhizopus rot Sour rot	8-16 fl oz/100 gal	Mix 8-16 fl oz of Chairman in 100 gallons of water or an appropriate water, wax/emulsion, or aqueous dilution of wax/oil emulsion.  Dip for approximately 30 seconds and allow fruit to drain.
High-Volume (Dilute Spray) Application	Black mold Gray mold Rhizopus rot Sour rot	8-16 fl oz/50,000 lb of fruit	Ensure proper coverage of the fruit.  Mix 8-16 fl oz of Chairman in an appropriate water, wax/emulsion, or aqueous dilution of a wax/oil emulsion for the crop treated.  Use T-jet or similar application system.
<b>Restrictions:</b>			
<b>Do not make more than one post-harvest application to the fruit.</b>			
<b>Do not apply more than 16 fl oz per 50,000 lb of fruit to tomato- small fruiting varieties.</b>			
<ul style="list-style-type: none"> <li>• Ensure the Chairman solution remains in suspension by using agitation.</li> <li>• Not for processing tomato.</li> </ul>			

Chairman™ and the Syngenta logo are trademarks of a Syngenta Group Company.