

WARARRANTY

To the extent consistent with applicable law, **Suterra® LLC** warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose stated in the Directions for Use. To the extent consistent with applicable law, **Suterra LLC** neither makes nor authorizes any agent or representative to make any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material. To the extent consistent with applicable law, **Suterra LLC's** maximum liability for breach of this warranty shall not exceed the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage, and use of this material contrary to label instructions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye and skin irritation. Avoid breathing vapor. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear chemical resistant gloves and protective eyewear.

FIRST AID STATEMENT

IF INHALED: Move person to fresh air. If person is not breathing call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for advice.

IF IN EYES: Hold eye(s) open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and continue rinsing. Call a Poison Control Center or doctor for advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER: Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. For emergency information on product, uses, and treatment, etc., call the National Pesticides Information Center at 1-800-858-7378, 6:30 am - 4:30 pm Pacific Time (PT), 7 days a week. Otherwise, call the Poison Control Center at 1-800-222-1222.

ENVIRONMENTAL HAZARDS

FOR TERRESTRIAL USES: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

CheckMate®

Puffer® CM-OFM

For the Control of Codling Moth and Oriental Fruit Moth in Apple, Pear, Peach, Nectarine and Any Other Crops Where the Codling Moth and Oriental Fruit Moth are Overlapping Problems.



FOR ORGANIC PRODUCTION

CheckMate® Puffer® CM-OFM contains a behavior modifying biochemical (pheromones) that disrupts the mating behavior of codling moth, *Cydia pomonella* and oriental fruit moth, *Grapholitha molesta*. Use in apple, pear, peach, nectarine and other orchards where the codling moth and oriental fruit moth are overlapping pests. **CheckMate Puffer CM-OFM** is used with the **CheckMate Puffer Aerosol Cabinet**, an automatic metered dispenser, to disrupt the nocturnal mating behavior of codling moth and oriental fruit moth.

ACTIVE INGREDIENTS*:

8,10 Dodecadien-1-ol, (E,E)	18.05 %
(Z)-8-Dodecen-1-yl acetate	11.63 %
(E)-8-Dodecen-1-yl acetate	0.75 %
(Z)-8-Dodecen-1-ol	0.12 %
Other Ingredients	69.45 %
Total	100.00 %

*Contains 93.85 grams total a.i. per canister

NET CONTENTS: 307 GRAMS (10.8 oz)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for First Aid and Precautionary Statements.

Suterra®

MANUFACTURED BY:

Suterra LLC
20950 NE Talus Place
Bend, Oregon, USA 97701
1-866-326-6737
Fax (541) 388-3705

EPA Registration No. 73479-11
EPA Establishment No. 56336-OR-1
BATCH CODE: See bottom of can.

© 2017 Suterra LLC. All Rights Reserved. CHECKMATE, PUFFER, AND SUTERRA are registered trademarks Suterra LLC.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING
Do not apply more than 150 grams of active ingredient per acre per year

METHOD OF APPLICATION

For use on apple, pear, peach, nectarine and other crops where the codling moth and oriental fruit moth are pests. In the spring, start applications prior to the sustained emergence (biofix) of oriental fruit moth adults and continue throughout the crop's susceptible period. Use a **CheckMate Puffer Aerosol Cabinet** to automatically dispense the contents of the **CheckMate CM-OFM** canister (herein referred to as "Puffer"). Placing cabinets in orchards and replacing canisters are to be done when cabinets are inactive, typically during daylight hours. To avoid accidental spraying, ensure that canister nozzles are pointed away during the performance of these tasks.

PRODUCT PLACEMENT

Suspend an activated **Puffer Aerosol Cabinet** containing a full **CheckMate Puffer CM-OFM** canister from a limb as high as possible in the tree canopy, ideally less than 3 feet from the top of the tree. The Puffers can also be placed slightly above the top of the canopy using posts or poles. Account for in-season growth of the canopy when placing. Place Puffer where spray is directed away from foliage and fruits. For best results, use in large contiguous plots. Distribute the Puffers throughout the orchard starting from the edges to achieve a rate between 1 and 2 Puffers per acre. In situations with a prevailing evening wind direction reinforce the upwind edge of the orchard. For irregularly shaped orchards consult your Suterra Technical Sales Representative.

APPLICATION NOTES

For best results apply at the time or shortly before codling moths or oriental fruit moths begin to emerge in the spring. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix (the first date moths are found consistently in monitoring traps) or during the growing season, the field must be treated with insecticides that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection. These supplemental insecticide applications must be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control young larvae. Monitor insect infestation with traps and by visual inspection of plants. Re-apply as needed, based on monitoring results and field scouting.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Store container in cool, dry area away from heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER HANDLING: Do not puncture or incinerate. Nonrefillable container. Do not reuse or refill this container. If can is empty: Remove from cabinet, place in trash or offer for recycling. If partly filled: Call your local solid waste agency for disposal instructions.



L17017 CM-OFM Std US FG16597 Rev07312018v1