

Bayer Environmental Science A Division of Bayer CropScience, LP 5000 CentreGreen Way, Suite 400 Cary, NC 27513 1-866-99BAYER (1-866-992-2937)

Cimarron Plus® Herbicide

2,4-D Tank Combinations for Control of Yucca

EPA Reg. No. 432-1572

Expiration: December 31, 2024

Product Bulletin

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Control of Yucca in Pasture and Rangeland

Apply Bayer CropScience Cimarron Plus® Herbicide in combination with 2, 4-D Low Volatile (LV) ester in the spring through the fall prior to frost. Make initial application from flower stalk elongation through seed pod development. Low volatile 2,4-D LV ester must be tank mixed for yucca and other companion species.

Initial Application Use Rates

milian / ippilication Goo mates	
Tank Mix	Rates Needed for Control
Cimarron Plus® Herbicide	0.625 to 1.25 oz pr/ac + 1.0 to 1.25 lb ai/ac
+ 2, 4-D LVE	

A second, follow-up application is recommended within two years of the initial treatment to control yucca seedlings and regrowth from rootstocks

Tank Mix	Rates Needed for Control
Cimarron Plus® Herbicide	0.375 to 0.625 oz pr/ac + 1.0 lb ai/ac
+ 2, 4-D LVE	·

Spray Adjuvants

Crop oil concentrate (COC), Modified Seed Oil (MSO) or Modified Seed Oil/Organosilicone (MSO/OS) are the preferred adjuvants. Aerial is the preferred application method. A minimum 4 gallons per acre volume is recommended for dense yucca populations.

Addition of ammonium nitrogen fertilizer may improve weed control. Use up to 2% v/v of a high quality urea ammonium nitrate (UAN), such as 28% N or 32% N, or up to 17 lb/acre of a spray grade ammonium sulfate (AMS).

Prior to application, contact the local Bayer representative for specific recommendations, application timing and adjuvants.

Variation in weather (moisture and temperature extremes), yucca physiological condition, soil type, and extent of yucca root system will determine treatment effectiveness

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

As with any herbicide product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).