

Source to Source

DIRECTIONS FOR USE

SOIL APPLICATIONS: Soil apply via mechanical application through direct spray or injection to the soil (banded, soil injection). Mix in clean tank with sufficient water for adequate mixing and coverage of target areas. It is not recommended in applications where material will come into direct contact with seeds or young roots in its concentrated form.



Foliar & Soil Applied Nutrient for California Trees, and Vines

0-0-0

Guaranteed Analysis

- Available Phosphate (P2O5).....5.0%
Cobalt (Co).....1.0%
Molybdenum (Mo).....3.0%
Derived From: phosphoric acid, cobalt chloride, sodium molybdate

DAANGER



HAZARD STATEMENT Causes severe skin burns and eye damage. Harmful if swallowed. Do not breathe dusts/mist/vapors. Wear protective gloves/protective clothing/eye protection/face protection. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product.

PRECAUTION STATEMENT Call a POISON CENTER or doctor/physician if you feel unwell. Rinse Mouth REFERENCE: SDS0013.02

GENERAL INFORMATION: CoMol™ 513 should be applied when practical experience, visual observation, or tissue or soil analysis indicate a need for Cobalt and Molybdenum. Cobalt & Molybdenum are important in reducing plant stress, stimulating root growth, and Nitrogen utilization. This product is for use in foliar sprays and soil applications. It is intended to be a supplement to a fertility program. Always monitor crop nutrient needs and responses with a soil, leaf tissue and irrigation water analysis program.

SEE LABEL ON PRODUCT FOR NET CONTENTS.

This material is available in: 2.5 GALLONS (9.46 LITERS) Density: 9.82 lbs per gallon (1.18 kgs per liter) @ 68°F



Manufactured by: Bio Huma Netics, Inc. 201 S. Roosevelt Ave. Chandler, AZ 85226

NOT FOR USE IN ORGANIC CROP AND ORGANIC FOOD PRODUCTION

Source to Source

Guaranteed & Distributed by: Source to Source, a TAP Family Company 3233 South "H" Street Tulare, CA 93274 Toll Free: 866.727.4572

FOLIAR APPLICATIONS:

CoMol 513 may be foliar applied. In a clean spray tank, add any dry powders and water and agitate. Add liquid Fertilizers and CoMol 513 last, continuing agitation. Concentration should not exceed 0.01% (volume to volume) in the finished spray solution. Always use enough water for uniform application and to prevent possible plant tissue damage to the intended crop from concentrated droplets. Contact your Source to Source representative for rates when using low-volume and electrostatic sprayers. Apply during the cool part of the day, or when temperatures below 85° F. Do not apply under windy conditions, or other conditions that promote fast drying of leaf surfaces.

Example, using maximum concentration of 0.01% =

Table with 3 columns: Product, in, Water. Rows: 19 oz in 15 gallons, 26 oz in 20 gallons, 36 oz\* in 50 gallons, 36 oz\* in 100 gallons, 36 oz\* in 150 gallons

\*36 oz is the maximum suggested rate for this material. Review Suggested Rates and Timings before application.

PRECAUTIONS: Agitate prior to use.

COMPATIBILITY: CoMol 513 is compatible with many acid-based nutrients. For unfamiliar mixes, an appropriate amount of the final mix should be applied to a small area of the intended crop to ensure plant compatibility prior to the planned application. User assumes full responsibility to ensure product, equipment and plant compatibilities.

STORAGE AND DISPOSAL: Store in a sealed container in cool, dry location, at temperatures above freezing. Triple rinse and dispose of containers according to local and federal regulations.

SUGGESTED RATES AND TIMINGS

ALMONDS, PISTACHIOS, WALNUTS & OTHER NUT CROPS

Apply in pre-bloom at 8-16 ounces per acre and post-bloom sprays to actively growing foliage at 8-32 ounces per acre. Repeat application in 14-21 days as needed. Water run 8-32 ounces per acre. Repeat in 21-28 days as needed.

DECIDUOUS FRUITS (Such as apples, cherries, nectarines, pears, peaches, plums & prunes)

Apply in pre-bloom at 8-16 ounces per acre and post-bloom sprays to actively growing foliage up to "pit hardening" at 8-32 ounces per acre. Repeat application in 14-21 days as needed. Water run 8-32 ounces per acre. Repeat in 21-28 days as needed. Post harvest application can enhance tree health and increase micronutrient levels in shoots and spurs for the following season.

CITRUS and AVOCADOS

Apply in pre-bloom and post-bloom sprays at 8-32 ounces per acre. Repeat application in 14-21 days as needed. Water run 8-32 ounces per acre. Repeat in 21-28 days as needed.

GRAPES and KIWI

Apply starting from bud break through 8-10" cane growth and continue at 14-21 day intervals through pre-bloom at rates of 8-16 ounces per acre. Apply again post-bloom through veraison as needed at 8-16 ounces per acre. Water run 8-32 ounces per acre. Repeat in 21-28 days as needed.

WARNING: Application of this material in excess may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. Buyer shall assume all risks of use including, but not limited to crop injury, ineffectiveness or other unintended consequences as may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the seller. In no case shall Source to Source be liable for consequential, special or indirect damages resulting from the use or handling of this product. Always refer to the label on the product before use. CoMol is Trade Mark of Source to Source, Tulare, CA. For information regarding the contents and levels of metals in this product, call your Source to Source Representative at (866) 727-4572. This label is for use in California and Arizona. For use in other states please contact your Source to Source dealer, as labels may vary by state.