



COLT + SWORD[®]

HERBICIDE

For selective postemergence control of annual and perennial broadleaf weeds and volunteer potatoes in small grains (barley, oats, wheat) and fallow cropland, and on-farm non-cropland.

ACTIVE INGREDIENT(S):

Fluroxypyr 1-methylheptyl ester: (4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic acid, 1-methylheptyl ester ¹	12.0%
2-methyl-4-chlorophenoxyacetic acid, 2-ethylhexyl ester ²	52.0%
OTHER INGREDIENT(S)	36.0%
TOTAL	100.0%

Contains xylene range aromatic solvent.

¹ Acid Equivalent: fluroxypyr: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic acid - 8.3% - 0.71 lb/gal
² Acid Equivalent: MCPA: 2-methyl-4-chlorophenoxyacetic acid - 33.3% - 2.84 lb/gal Isomer Specific AOAC Method

KEEP OUT OF REACH OF CHILDREN CAUTION—PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565. Note to Physician: May pose an aspiration pneumonia hazard. Contains xylene range aromatic solvent.</p>	

EPA REG. NO. 34704-1011

EPA EST. NO. 34704-MT-001

NET CONTENTS 2.5 GAL (9.46 L)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below.

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton when mixing, loading, or using any hand-held equipment.
- Shoes plus socks

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland.

These mixers/loaders also must wear:

- Chemical-resistant apron
- NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N², R, P, or HE filter.

See Engineering Controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protections Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and may be toxic to aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

MCPA has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms or nurseries: **When this product is applied to non-cropland areas, keep unprotected persons out of treated areas until sprays have dried.**

PRODUCT INFORMATION

Colt®+Sword® Herbicide is a selective postemergence product for control of annual and perennial broadleaf weeds and volunteer potatoes in wheat, barley, or oats not under seeded with a legume, fallow cropland and on-farm non-cropland uses such as fence rows, building perimeters, around irrigation equipment and roadways.

Application Precautions and Restrictions

- Do not apply this product directly to, or otherwise permit it to come in direct contact with, susceptible crops or broadleaf plants including alfalfa, cotton, lettuce, edible beans, lentils, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, tobacco, grapes, legumes, fruit trees, canola, tame mustard, other vegetables or ornamentals.

Vapors from this product may injure susceptible plants in the immediate vicinity.

- Avoid applications where proximity of susceptible crops or other susceptible broadleaf plants is likely to result in exposure to spray or spray drift.
- Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not apply in greenhouses.
- **Maximum Application Rate:**
 - Do not apply more than 2.0 pints of Colt+Sword Herbicide (0.178 pound acid equivalent floxypyr plus 0.71 pound acid equivalent MCPA) per acre per growing season.
- **Plant-back Restriction:**
 - Plant only those crops listed on this label or Federally approved supplemental labeling for Colt+Sword Herbicide within 120 days following application. Do not plant rotational crops within 120 days of application to fallow croplands.
- **Chemigation:** Do not apply this product through any type of irrigation system.

Management of Kochia Biotypes: Research has suggested that many biotypes of Kochia can occur within a single field. While Kochia biotypes can vary in their susceptibility to Colt+Sword Herbicide, all will be suppressed or controlled by the 1.5 pints labeled rate. Application of Colt+Sword Herbicide at rates below the 1.5 pints rate can result in a shift to more tolerant biotypes within a field.

Best Resistance Management Practice: Extensive populations of dicamba tolerant Kochia have been identified in certain small grain and corn production regions (such as Chouteau, Fergus, Liberty, Toole, and Treasure counties in the state of Montana). In these areas, Colt+Sword Herbicide should be applied at a minimum rate of 1.5 pints per acre for optimal control of dicamba tolerant Kochia. In addition, Colt+Sword Herbicide should be rotated with products that do not contain dicamba to minimize selection pressure. Use of these practices will preserve the utility of Colt+Sword Herbicide for control of dicamba tolerant Kochia biotypes.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when wind speed is 2 to 10 mph at the application site.

Aerial Application:

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
2. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.
3. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Do not make applications into temperature inversions.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information:**

Ground Applications: To minimize spray drift, apply Colt+Sword in a total spray volume of 8.0 or more gallons per acre using spray equipment designed to produce large-droplet, low pressure sprays. Refer to the spray equipment manufacturer's recommendations for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Spot treatments should be applied only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Importance of Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size:

Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure-Use the lower spray pressures for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Do not make applications into temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Sprayer Cleanup

To avoid injury to or exposure of nontarget crops, thoroughly clean and drain spray equipment used to apply Colt+Sword Herbicide after use. Cleaning should occur as soon as possible after application of Colt+Sword Herbicide. Spray equipment should be cleaned after use with Colt by the following procedure:

1. Drain any remaining Colt+Sword Herbicide from the spray tank and dispose of according to label disposal instructions.
2. Hose down the interior surfaces of the tank. Flush tank, hoses, boom, and nozzles with clean water for 10 minutes. Fill the tank with water and recirculate for 15 minutes. Spray part of the mixture through the hoses, boom, and nozzles and drain the tank. All rinse water must be disposed of in compliance with local, state, and federal guidelines.
3. Remove the nozzles and screens and clean separately.
4. If the spray equipment will be used on crops other than those labeled for Colt+Sword Herbicide, repeat steps 1 and 2 and thoroughly wash the outside of spray tank and the boom.

MIXING INSTRUCTIONS

Colt+Sword Herbicide

Fill the spray tank approximately 1/2 to 3/4 full with water. Add the required amount of Colt+Sword Herbicide, and then finish filling the spray tank. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixing

Colt+Sword Herbicide may be applied in tank mix combination with labeled rates of other herbicides provided (1) the tank mix product is labeled for the use site (timing and method of application is the same as Colt+Sword Herbicide); and (2) tank mixing with Colt+Sword Herbicide is not prohibited by the label of the tank mix product.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed application rates. If products containing the same active ingredient are tank mixed, do not exceed the maximum allowable active ingredient use rates.
- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: Perform a jar test prior to tank mixing to ensure compatibility of Colt+Sword Herbicide and other pesticides, fertilizers, or carriers. Use a clear glass jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Tank Mixing Instructions:

Fill the spray tank to approximately 1/4 to 1/3 of the total spray volume required. Start agitation. Add different formulation types in the order indicated, allowing time for complete mixing and dispersion after addition of each.

1. Add dry flowables; wettable powders; aqueous suspensions, flowables or liquids.
2. Maintain agitation and fill spray tank to 3/4 of total spray volume and then add Colt+Sword Herbicide and other emulsifiable concentrates and any solutions.

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

APPLICATION DIRECTIONS

Application Timing: Apply to actively growing weeds. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Only weeds that are emerged at the time of application will be affected.** Foliage that is wet at the time of application may decrease control.

Colt+Sword Herbicide applications are rain-fast within 1 hour after application.

Application Rates: Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds) the higher rates within the rate range will be needed. Weeds growing in the absence of crop competition generally require higher rates to obtain satisfactory control or suppression.

Effect of Temperature on Herbicidal Activity: Herbicidal activity of Colt+Sword Herbicide is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum herbicidal activity is 55 °F to 75 °F. Reduced activity will occur when temperatures are below 45 °F or above 85 °F. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance.

Coverage: For best results, apply in 3.0 or more gallons per acre by air or in 8.0 or more gallons per acre by ground equipment. Do not exceed 40.0 gallons per acre total spray volume. Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Inadequate spray volume and coverage may result in decreased weed control. As crop canopy and weed density increase, spray volume should be increased to obtain equivalent weed control. Use larger nozzle tips or decrease spraying speed to increase spray volume rather than increasing boom pressure. Refer to manufacturer's recommendations for information on relationships between spray volume, and nozzle size and arrangement.

Adjuvants: Use of a high quality adjuvant labeled for use on growing crops is suggested for improved weed control. Adjuvants are especially beneficial when applications are made (a) at lower carrier volumes, (b) under conditions of cool temperature, low relative humidity or drought, or (c) to small, heavily pubescent Kochia.

Spot Treatments: To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to directions provided below.

Hand-Held Sprayers: Hand-held or backpack sprayers may be used for spot applications of Colt+Sword Herbicide if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1000 square feet. Mix the amount of Colt+Sword Herbicide (fluid ounce or milliliter) corresponding to the desired broadcast rate in 1.0 or more gallons of spray. To calculate the amount of product required for larger areas, multiply the table value (fluid ounce or milliliter) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3500 square feet, multiply the table value by 3.5 (calc. 3500 ÷ 1000 = 3.5). An area of 1000 square feet is approximately 10.5 X 10.5 yards (strides) in size.

**Amount of Colt+Sword Herbicide to Equal Specified Broadcast Rate
(Mix with 1.0 Gallon or More of Water and Apply to 1000 Square Feet)**

	1.5 Pt/A	2.0 Pt/A
	0.55 fl oz (16.0 ml)	0.73 fl oz (22.0 ml)

1.0 fl oz = 29.6 (30.0) ml

WEEDS CONTROLLED OR SUPPRESSED

(Numbers in parentheses (-) in weeds list refer to footnotes below.)

Weeds Controlled

Bedstraw (cleavers)	Dandelion	Marshelder	Ragweed, common
Bittercress	Dock, curly	Morningglory, annual	Ragweed, giant
Bull nettle	Flax, volunteer	Mousetail	Russian thistle
Bur beakchervil	Flixweed	Mustard (except Blue) (2)	Shepherd's-purse
Buttercup	Geranium, Carolina	Pennycress, field	Sicklepod
Canola, volunteer	Hemp dogbane	Pigweed	Sowthistle, annual
Chickweed	Horseweed	Plantains	Sowthistle, spring
Cinquefoil	Jacob's ladder	Prickly lettuce	Speedwell
Cocklebur	Kochia (1)	Primrose, evening	Sunflower
Coffeeweed	Lambsquarters, common	Puncturevine	Sweetclover
Copperleaf, Virginia	Mallow, Venice	Purslane, common	Velvetleaf

1. Includes herbicide tolerant biotypes.
2. Apply prior to bolting.

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

Weeds Suppressed (1)

Alfalfa	Croton	Jimsonweed	Redstem filaree
Beggarticks	Daisy	Knawel	Smallseed falseflax
Bindweed, field	Dragonhead mint	Ladysthumb	Smartweed
Buckwheat, wild	Fiddleneck	Mallow, common	Tansymustard
Burcucumber	Field bindweed	Nightshade species	Thistle, bull
Burdock, common	Galinsoga	Onion, wild	Thistle, Canada
Carpetweed	Garlic, wild	Peppergrass	Thistle, musk
Carrot, wild	Goatshead	Pepperweeds (annual)	Vetch, hairy
Catsear	Hempnettle	Poison hemlock	Yellow rocket
Clover, red	Henbit	Potato, volunteer	
Corn spurry	Hoarycross	Purslane, common	
Cowcockle	Ironweed	Radish, wild	

1. **Suppression** is expressed as a reduction in weed competition (reduction population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

APPLICATION SITES

Crop Uses

Wheat (Including Durum), Barley, and Oats

Apply as a broadcast postemergence treatment to actively growing wheat (including durum), barley or oats, from the 3-leaf crop growth stage up to and including flag leaf emergence (Zadoks scale 39) for control of broadleaf weeds. Apply when weeds are actively growing, but before weeds are 8 inches tall or vining. For control of Volunteer potatoes, apply before potato plants are 8 inches tall. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperatures prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Do not use if cereal crop is underseeded with a legume.**

Spot Application: Spot applications may be made, however, to prevent over-application spot treatments should be applied at rates and spray volumes equivalent to broadcast application. See instructions for "Spot Application" in "Application Directions" section.

Broadcast Application Rates:

(Numbers in parentheses (-) refer to footnotes following table.)

Weed Size or Species ^a	Application Rate (Pt/A)
Susceptible broadleaf weed seedlings less than 4 inches tall ^b	1.125
Susceptible broadleaf weed seedlings less than 8 inches tall or vining	1.5
Volunteer potatoes	1.5 to 2.0 ^c

^a See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

^b The 1.125 pints per acre rate will generally provide satisfactory control of Kochia seedlings less than 4 inches tall (including ALS resistant biotypes). However, when conditions for control are less favorable, such as under drought or cool temperatures, the 1.5 pints per acre rate will provide more consistent control of Kochia seedlings 1 to 4 inches tall. Control of small Kochia with reduced rates will be more consistent if Kochia is at least 1 inch tall. The 1.5 pints per acre rate should be used for optimal control of dicamba tolerant Kochia populations (see "Management of Kochia Biotypes" in the General Information section of this label).

^c Crop injury may occur at rates higher than 1.5 pints per acre.

Restrictions:

- Do not spray plants beyond the flag leaf emergence stage.
- Do not allow livestock to graze treated areas within 45 days of application.
- Do not apply more than 2.0 pints (0.178 pound acid equivalent fluroxypyr plus 0.71 pound acid equivalent MCPA) per acre of Colt + Sword.
- Do not make more than 1 application of this product per season.
- Do not allow dairy animals or meat animals being finished for slaughter to forage or graze treated areas within 7 days after application.
- **Preharvest Interval:** Do not apply within 14 days before cutting of hay or 40 days before harvesting of grain and straw.

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

Fallow Cropland

Apply as a single broadcast treatment by ground or aerial equipment to control susceptible broadleaf weeds. Apply when weeds are actively growing, but before Kochia is 8 inches tall and before Wild buckwheat is vining. Colt+Sword Herbicide may be applied at labeled rates alone or in tank mix combination with other herbicides (See tank mixing precautions in "Mixing Instructions" section.)

Restriction:

- A plant back interval of 120 days is required for fallow croplands.

Broadcast Application Rates:

Weed Size or Species†	Application Rate (Pt/A)
Susceptible broadleaf weed seedlings less than 8 inches tall or vining	1.5 to 2.75
Volunteer potatoes	1.5 to 2.75

† See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

On-Farm Non-Cropland

Apply as a single broadcast treatment or spot treatment to control susceptible broadleaf weeds in on-farm non-cropland areas such as fence rows, building perimeters, around irrigation equipment and on-farm private roadways. Apply at the rate of 1.5 to 2.75 pints per acre when weeds are small and actively growing, but before weeds are 8 inches tall or vining. Spot treatments should be applied at rates and spray volumes equivalent to broadcast application. See instructions for "Spot Application" in "Application Directions" section. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

CRP Acres

Do not use on CRP acres that are underseeded with desirable legumes, clovers, or other sensitive broadleaf plants.

Colt+Sword Herbicide may be applied to Conservation Reserve Program (CRP) acres. For best results, apply as a single broadcast treatment by ground or aerial equipment to control susceptible broadleaf weeds. Apply at the rate of 1.5 to 2.75 pints per acre when weeds are small and actively growing, but before weeds are 8 inches tall or vining. Spot treatments should be applied at rates and spray volumes equivalent to broadcast application. See instructions for "Spot Application" in "Application Directions" section. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

Restriction: Grazing or haying of treated CRP acres is prohibited.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or fertilizer by storage or disposal.

PESTICIDE STORAGE: Store above 10 °F or warm and agitate before use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 50 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank

COLT®+SWORD® HERBICIDE
EPA REG. NO. 34704-1011

Storage & Disposal cont'd.:

or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 50 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, READ THE ENTIRE DIRECTIONS FOR USE AND THE FOLLOWING CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY. BY BUYING OR USING THIS PRODUCT, THE BUYER OR USER ACCEPTS THE FOLLOWING CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY, WHICH NO EMPLOYEE OR AGENT OF LOVELAND PRODUCTS, INC. OR THE SELLER IS AUTHORIZED TO VARY IN ANY WAY.

FOLLOW THE DIRECTIONS FOR USE OF THIS PRODUCT CAREFULLY. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP OR OTHER PLANT INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT FROM SUCH RISKS AS WEATHER OR CROP CONDITIONS, MIXTURE WITH OTHER CHEMICALS NOT SPECIFICALLY IDENTIFIED IN THIS PRODUCT'S LABEL, OR USE OF THIS PRODUCT CONTRARY TO THE LABEL INSTRUCTIONS, ALL OF WHICH ARE BEYOND THE CONTROL OF LOVELAND PRODUCTS, INC. AND THE SELLER. THE BUYER OR USER OF THIS PRODUCT ASSUMES ALL SUCH INHERENT RISKS.

SUBJECT TO THE FOREGOING INHERENT RISKS, LOVELAND PRODUCTS, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE PURPOSES STATED IN THE DIRECTIONS FOR USE WHEN THE PRODUCT IS USED IN STRICT ACCORDANCE WITH SUCH DIRECTIONS FOR USE UNDER NORMAL CONDITIONS OF USE. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Colt and Sword are registered trademarks of Loveland Products, Inc.

FORMULATED FOR
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286