Copper 5%



Guaranteed Analysis

Copper (Cu)5.0% 5.0% water soluble Copper (Cu)

Derived from copper sulfate.

F002633

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

BACOR® products are compatible with most refilizers, insecticides, fungicides and herbicides (follow product labels) which may incorp and the product should be useful to the product should be lested by a licensed and experienced applicator with such mixtures (jar test) and small test plots applied on soil/crop before any large scale for testiments are started.

ONLY A LICENSED AND EXPERIENCED APPLICATOR SHOULD APPLY THIS PRODUCT.

Limited Warranty Manufacturer warrants that this product conforms to the guaranteed analysis for the growing season in which purchased. Manufacturer makes no other express or implied warranty of fitness or merchantability or any other express or implied warranties, in no such case shall manufacturer or the seller be liable for consequential, incidental, special, exemplary or indirect damages resulting from the use or handling of this product, any and all damages claimed shall be limited to the amount caid for the product.

Manufacturer offers this product and the Buyer/User accepts it subject to the forgoing Conditions of Sale and Limited Warranty

WARNING: This product may stain (Keep off unwanted areas such as, sidewalks, patios, driveways, fences, painted areas). As a precaution: Spray off unwanted areas with only water after applying.

Hazard Statement May be harmful if swallowed. Harmful in contact with skin. Causes skin irritation. Causes serious eye irritation. Harmful if inhaled. Harmful to aquatic life. Harmful to aquatic life with long lasting effects.

Precautionary Statement

Prevention: Avoid breathing vapors. Wash thoroughly after handling. Use outdoors or in a well-ventilated area. Avoid release to the environment. Wear protective clothing. Wear eye protection.

Response: If on skin: Wash with plerity of wafer. If Inhaled: Remove person for fresh air and keep comfortable for breathing. If in eyes: Rinse cations with water for several minutes. Bennevo contact lenses, if present and easy to do. Continue risining, Call a poison center if you feel unwell. If skin irritation occurs: Get medical advice. If eye irritation persists: Get medical attention. Take off contaminated coldining and wash before reuse.

Storage: Store away from incompatible materials. Products should be stored above 12.8°C (55°F).

Storage of bulk products is best suited for original manufacturing containers. If required, storage in a polypropylene storage tank with polypropylene fittings is recommended. Metal storage tanks and fittings are not recommended for long term. Disposat: Dispose of contents in accordance with local regulations.

Plant Nutrients To Prevent & Correct Deficiencies For Soil & Foliar Application APPLICATION RATES & DIRECTIONS: Unless advised by consultant, do not apoly during bloom

APPLICATION RATES & DIRECTIONS: Unless advised by consultant, do not apply during bloon stage. Recommendations listed are only general. Plant species and varieties may have wide variations in their sensitivity or requirement to foliar/soil applications of certain nutrients. Optimum rate of application and frequency will vary between fields, depending on crop, soil pH, organic matter content, etc. Test application (test plot) to assess the potential for undesirable effects under specific conditions is essential.

A surfactant (detergent) is recommended for all foliar applications to decrease drop size and enhance wetting of entire leaf surface. For best results apply as a foliar spray early in the morning or late evening. Do not apply during hot, dry weather, or in periods of drought or when plants are under stress.

ADD WATER FIRST TO TANK OR SPRAYER BEFORE ADDING PRODUCT!

AERIAL APPLICATIONS: Use at least 20 parts water to 1 part of BAICOR® Copper 5%. Add at least 20 parts water before introducing product.

FOLIAR APPLICATIONS: Use at least 100 parts water to 1 part BAICOR® Copper 5%. Add at least 100 parts water before introducing product.

FRUIT, NUT & VINE CROPS: Including (but not limited to) almonds, hazelnuts, grapes, pecans and walnuts. Apply 4 - 8 ounces per acre. NOTE: Before applying to pome or stone fruit, consult your qualified and licensed consultant for recommendations.

FIELD AND VEGETABLE CROPS: Apply 4 - 8 ounces per acre.

GRAIN CROPS: Apply 4 to 8 ounces per acre at 3-4 leaf stage.

TURF GRASSES: Apply 4 to 8 ounces per acre.

 $\label{eq:sprinkler} \textbf{SPRINKLER IRRIGATION:} \ \ \text{Apply } 16-32\ \text{oz's}\ (0.5-1\ \text{qt})\ \text{per acre with irrigation water.} \ \ \text{Use check valve to prevent back flow into water system.}$

SOIL APPLICATION RATES: Use at least 20 parts water to 1 part BAICOR® Copper 5%. Do not apply directly to seeds unless it has been determined/tested by the consultant or grower that it is not harmful or injurious to the seed.

Supplemental Information

This product is a "Hazardous Chemical" as defined by the OSHA Hazard Communication Standard, 29 CFB 1910 1200

All components are on the U.S. FPA TSCA Inventory List.



CAUTION: KEEP OUT OF REACH OF CHILDREN

10.12 lbs/gal @ 68° F

REV 9/10/19

Lot #

For more information please visit our website:

2.5 Gallons/9.46 L 25.3 lbs./11.5 kg 5 Gallons/18.93 L 50.6 lbs./23 kg 55 Gallons/208.2 L 556.6 lbs./253 kg 270 Gallons/1022 L 2,732 lbs./1,239 kg