



NOTICE

Before using this product, read the entire label

Coptrac™

YaraVita™

4-0-0

Coptrac™

A formulated product for the treatment of copper deficiency by foliar or soil application

Guaranteed analysis:

Total Nitrogen(N).....4%

4% Urea Nitrogen

Copper(Cu).....33%

Derived from Urea, Copper (I) Oxide

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>


This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

**2½ gallons/
9.4 litres**

F1673

**Net weight
31 lb 12oz/
14.4 kg**

Contents made in Great Britain
Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road,
The Industrial Estate, Pocklington, York YO42 1DN, U.K.
Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200,
Tampa, FL 33602 Customer Service: 800-234-9376




YaraVita Coptrac™

THIS PRODUCT IN NON-BULK PACKAGING IS A MARINE POLLUTANT WHEN TRANSPORTED BY VESSEL. HOWEVER, IT IS NOT DOT REGULATED WHEN TRANSPORTED BY HIGHWAY ONLY PER 49 CFR, 171.4(c)

UN 3082

Environmentally hazardous substance, liquid, n.o.s. (dicopper oxide)



Danger

H302: Harmful if swallowed.(kidneys)

H318: Causes serious eye damage.

H373: May cause damage to organs through prolonged or repeated exposure.

H410: Very toxic to aquatic life with long lasting effects.

P280: Wear protective gloves and eye protection.

P270: Do not eat, drink or smoke when using this product.

P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

P301+P312: IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.

P391: Collect spillage.

YaraVita™ Coptrac™

APPLICATION RATES AND TIMINGS

Alfalfa: 0.5 pint/acre applied at start of tillering.

Water rate: 20 gallons/acre.

Carrots: ½ pint/acre when crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.

Cereals: ½ pint/acre from 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). For fall-planted crops, an application prior to winter dormancy is strongly recommended. For moderate to severe deficiency, repeat the application once or twice at 10-14 day intervals. Water rate: 20 gallons/acre.

Conifers: 2 applications of ½ pint/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

Corn: ½ pint/acre at 4 to 6 leaf stage.

Water rate: 20 gallons/acre.

Garlic: ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6 inches tall. Repeat applications at 10 to 14 day intervals.

Water rate: 7.5 to 20 gallons/acre.

Nursery/Ornamentals: One pint in 100 gallons water (0.125% v/v) as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Note: Do not apply within one month of picking/marketing.

Maximum water rate: 20 gallons/acre.

Onions: ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.

Potatoes: 0.5 pint/acre applied 7 to 14 days after 100% emergence. Water rate: 20 gallons/acre.

Rice: ½ pint/acre applied at start of tillering. Water rate: 20 gallons/acre.

Sorghum: ½ pint/acre at the 4 to 8 leaf stage. Water rate: 3 to 20 gallons/acre.

Sweet Corn: ½ pint/acre at 4-6 leaf stage. Water rate: 20 gallons/acre.

SOIL APPLICATION

All Crops: 2 quarts/acre to 1 gallon/acre, applied pre-planting or pre-emergence. Water rate: 5 gallons/acre minimum

Extended recommendations available at
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and reproductive Harm - www.P65Warnings.ca.gov