

	FIFRA 2(ee) Recommendation
	EPA Reg. No. 279-9606
	FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA
	THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for CORAGEN® insect control is valid until December 31, 2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR USE VIA DRIP CHEMIGATION APPLICATION IN GLOBE ARTICHOKE FOR CONTROL OF BEET ARMYWORM AND CABBAGE LOOPER

Directions For Use
<p>PESTS CONTROLLED CORAGEN insect control is recommended for the control of beet armyworm and cabbage looper via drip chemigation in globe artichoke.</p> <p>Read the chemigation section of the federal label for more information on how to use via drip chemigation.</p> <p>See the RESISTANCE MANAGEMENT section of the CORAGEN insect control Federal label for additional guidance regarding insecticide resistance management.</p>

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Artichoke, globe	Drip chemigation	Beet armyworm Cabbage looper	0.045 - 0.098	3.5 - 7.5	3	4
<p>USE RESTRICTIONS</p> <p>Begin drip application when insect populations first appear. CORAGEN insect control must be applied uniformly in the root zone or poor performance will result.</p> <p>Drip tape must be placed directly underneath a single row to ensure CORAGEN insect control is applied in the root zone.</p> <p>During the first half of the crop growing cycle, allow 1-3 days for CORAGEN insect control to be translocated into the leaves for pest protection. For applications made in later stages of plant development, translocation of CORAGEN insect control into aerial portions of the plant may take up to 7-10 days.</p> <p>Do not make more than 2 drip chemigation applications of CORAGEN insect control per crop.</p> <p>Minimum application interval between drip chemigation treatments is 14 days.</p> <p>Apply no more than 15.4 fl oz CORAGEN insect control or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.</p>						

R-4487 092820 10-09-20

FMC, the FMC logo and Coragen are trademarks of FMC Corporation or an affiliate.
© 2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104