

Below recommended rates in up to 100 gallons of water per acre.

CROP TYPE	PEST	RATE (Oz./Ac.)
Tomatoes, Potatoes Peppers	Anthraxnose (Colletotrichum coccodes), Black mold (Alternaria alternata), Buckeye rot (Phytophthora spp.), Early blight (Alternaria solani), Powdery mildew (Oidium sp. sica), Septoria leaf spot (Septoria lycopersici), Target spot (Corynespora Cassicola), Late blight (Phytophthora infestans)	8-16
Bassica leafy greens: Broccoli, Brussels sprouts, Chinese cabbage, Chinese mustard Cauliflower	Alternaria leaf spot (Alternaria spp.), Downy mildew (Peronospora parasitica), White rust (Albugo candida), Black spot (Alternaria spp.), Cercospora (Cercospora spp.), Seedling root rot, basal stem rot (Rhizoctonia solani)	8-16
Cucurbits: Cantalope, Cucumber, Honeydew Melons, Muskmelon, Watermelon, Pumpkin, Squash, Zucchini	Cucurbits: Cantaloupe, Cucumber, Anthracnose (Colletotrichum coccodes), Belly rot (Rhizoctonia solani), Downy mildew (Pseudoperonospora cubensis), Gummy stem blight (Didymella bryoniae), Leaf spots (Alternaria spp., Cercospora spp.), Myrothecium canker Myrothecium rostratum, Plectosporium blight (Plectosporium tabacinum), Powdery mildew (Sphaerotheca fuliginea, Erysiphe chichoracearum)	8-16
Turf and Ornamentals	Anthraxnose (Colletotrichum graminicola), Brown Patch (Rhizoctonia solani), Cool Weather Brown/Patch Yellow Patch (Rhizoctonia cerealis), Fairy Ring (Lycoperdon spp., Agrocybe pedicels, and Bovista plumbea), Fusarium Patch (Microdochium nivale), Gray Leaf Spot (Pyricularia grisea), Gray Snow Mold Typhula Blight (Typhula incarnata), Leaf Rust, Stem Rust, Stripe Rust (Puccinia spp.), Leaf Spot (Bipolaris sorokiniana)	8-16
Soybeans, Corn, Sweet Corn, Sorghum, Cotton, Tobacco	Aerial blight (Rhizoctonia solani), Anthracnose (Colletotrichum truncatum), Alternaria leaf spot (Alternaria spp.), Brown spot (Septoria glycines), Cercospora Blight and leaf spot (Cercospora kikuchii), Rust, Powdery mildew, Downy mildew, Frogeye leaf spot (Cercospora soja), Pod and stem blight (Diaporthe phaseolorum), Rust (Phakopsora spp.), Southern blight (Sclerotium rolfsii), Rhizoctonia solani Rhizoctonia solani Rust, Stem Rust, Stripe Rust (Puccinia spp.), Leaf Spot (Bipolaris sorokiniana)	8-16
Potatoes	Early blight (Alternaria solani), Late blight (Phytophthora infestans), Black dot (Colletotrichum coccodes), Powdery mildew (Erysiphe chichoracearum), Black scurf (Rhizoctonia solani), Silver scurf (Helminthosporium solani)	8-16
Strawberries, Blueberries Raspberries, Blackberries	Downy mildew (Peronospora destructor), Botrytis leaf blight (Botrytis aclada), Rhizoctonia damping-off (Rhizoctonia solani)	8-16
Pear	Fire blight, Blossom blast, (Pseudomonas spp.)	8-16
Bulb Vegetables: Garlic Leek Onion, Bulb Onion, green Welsh onion, shallot	Cladosporium leaf blotch (Cladosporium allii), Purple blotch (Alternaria porri), Rust (Puccinia allii), Downy mildew (Peronospora destructor), Botrytis leaf blight (Botrytis aclada), Rhizoctonia damping-off (Rhizoctonia solani)	8-16

Application Instructions: Do not apply more than two consecutive applications and rather rotate CYCLOPS® Organic Fungicide to an alternative chemistry for two applications. Apply early in the morning or in late afternoon. Spray directly onto the pathogen(s). Repeat applications at an interval of 4-7 days, depending on bacteria infestation pressure. To prevent product degradation, do not use spray water with a pH lower than 4.0 or higher than 8.0. Use pH correctors for pH's outside of these ranges.

HAND HELD EQUIPMENT: When using hand held equipment use 0.5 oz - 1.0 oz of CYCLOPS™ Organic Fungicide/ gallon of water.

Final mix volume	Fluid Oz of CYCLOPS®	Tbs of CYCLOPS®	Apply no more than once in a 7-day period. Repeat application as necessary. Coverage is essential to establish control. The use of a spreader/sticker may increase contact and efficacy of treatment. Check compatibility by spraying a small number of plants before initial application. Pre Harvest Interval: CYCLOPS® Organic Fungicide may be applied on crops listed on recommended use table at anytime and up to and including the day of harvest.
1 Gallon	0.06	0.5	
5 Gallons	0.30	2	
10 Gallons	0.60	4	
25 Gallons	1.50	10	

PHYTOTOXICITY: this product is not phototoxic and will not burn when used in accordance with label use instructions.