



Section 2(ee) Recommendation

Date: February 10, 2012

Product: Daconil Action™
EPA Reg. No. 100-1364

Use: Reduced Rates of Daconil Action for the control of dollar spot on turf

State: TX

DIRECTIONS FOR USE

Daconil Action fungicide couples a multiple-site contact fungicide with a systemic compound used for control and suppression of disease through induction of host plant resistance. Daconil Action exhibits an additional unique mode of action, beyond chlorothalonil alone, which mimics the natural systemic activated resistance (SAR) response found in most plant species. This SAR response does not involve direct activity against the target pathogens.

This recommendation applies to Group A Turf as listed on the federal label.

Golf Course Fairways and Roughs, professional and collegiate athletic field; and lawns around commercial and industrial buildings

Daconil Action provides control of dollar spot (*Sclerotinia homeocarpa*, *Lanzia*, or *Moellerodiscus* spp.) on turf when applied at the following preventative rates under low to moderate disease pressure. Under conditions of moderate to severe disease pressure, refer to use rate tables on federal label.

Pre-Disease Rates			
Disease Controlled	Application Interval (Days)	Use Rate (fl. oz. product per 1,000 sq. ft.)	Use Rate (Pints Product per Acre)
Dollar Spot	14	1.6 – 2.0	4.4 – 5.4

Note: These rates are not to be used if disease is present. Consult the federal label for post-disease rates.

The user must refer to the federally approved label and read and follow all directions for use, restrictions, and precautions

The user should have this recommendation in their possession at the time of use.

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the EPA.

Daconil Action™ and the Syngenta logo are trademarks of a Syngenta Group Company.

©2012 Syngenta Crop Protection, LLC,
410 Swing Road, Greensboro, NC 27409

TX1364097AC0212