



**FIFRA 2(ee)
RECOMMENDATION
Broadcast Application Methods
in California Onions**

FIFRA 2ee RECOMMENDATION

DACTHAL® Flowable Herbicide

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

EPA Reg. No.: 5481-487

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS BULLETIN MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Before applying **DACTHAL Flowable herbicide** please read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of **DACTHAL Flowable herbicide** according to this supplemental labeling is subject to the use precautions and limitations imposed by the label on the container.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered **DACTHAL Flowable herbicide** label before using this product.

Any Application of DACTHAL FLOWABLE Should Be Made Prior to Weed Seed Germination

Use Directions

Crop	When to Apply	Rate per acre	Remarks
Onions [dry bulb onions, dry bulb shallots and green onions (green onions, leeks, spring onions or scallions, Japanese bunching onions, green shallots or green eschalots)]	At seeding or transplanting and/or at layby.	See the table below for rate information. Mixtures of low rates of DACTHAL FLOWABLE with certain herbicides, including pendimethalin, can be effective in weed control.	Apply uniformly to the soil as a spray. DACTHAL FLOWABLE can be sprayed directly over transplants without injury. A layby application can be made on onions either alone or in addition to a DACTHAL FLOWABLE preemergence treatment (with a minimum of 14 days between treatments) up to 14 weeks after planting at rates up to 14 pints per acre on any soil type. If weeds emerge prior to layby, the onions should be cultivated or weeded prior to application. Use ¼ inch of overhead irrigation to incorporate DACTHAL FLOWABLE into the soil. Where onion emergence is expected to be slow due to cold and wet soil conditions, delay application until seeds begin to germinate to reduce the risk of crop injury. Preplant incorporation is not recommended for onions. Broadcast applications of DACTHAL FLOWABLE may be applied during the period from January 1 through July 31 in the following counties: Siskiyou, Modoc, Shasta and Lassen.

Rates of Application and Weeds Controlled: DACTHAL FLOWABLE is recommended at rates of 6-14 pints/A depending on soil type and weeds to be controlled. The following table outlines general recommendations for the use of DACTHAL FLOWABLE on various soil types. The maximum rate for each soil type should be used when moderately susceptible weeds are present. Consult the UC extension specialist or your AMVAC representative for specific recommendations. ***DACTHAL FLOWABLE should not be applied under conditions of cool, wet weather when plants are not actively growing.*** Refer to specific label recommendations for onions for further details.

Soil Type	Organic Matter %	Broadcast Rate of DACTHALFLOWABLE/A
Light - sand or sandy loam	0-3	6-8 pints
Medium - silt loam	0-3	8-10 pints
Heavy - clay loam	0-3	10-14 pints
Heavy - silt loam	3-5	12-14 pints

©2020 AMVAC Chemical Corporation. A wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. American Vanguard, AMVAC, the AMVAC logo, and DACTHAL are trademarks owned by AMVAC Chemical Corporation.

Registrant:

AMVAC Chemical Corporation
 4695 MacArthur Court, Suite 1200
 Newport Beach, CA U.S.A.
 1-888-462-6822