

FIFRA Section 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Delegate[®] WG

EPA Reg. No. 62719-541

EPA SLN No. OR-170016

(For Distribution and Use Only in the State of Oregon)

Suppression of Spotted Wing Drosophila and One-day Pre-harvest Interval in Blueberry

Expiration Date: This label is valid until December 31, 2022 or until otherwise amended, withdrawn, canceled, or suspended.

Group	5	INSECTICIDE
--------------	----------	--------------------

Active Ingredient:

spinetoram (a mixture of
spinetoram-J and spinetoram-L)25.0%

Other Ingredients75.0%

Total100.0%

Contains 25% active ingredient on a weight basis (250 g ai/kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

Environmental Hazards

This product is toxic to bees exposed to treatment during the 3 hours following treatment. Do not apply this pesticide to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on the label.

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Special Local Need (SLN) label and the EPA registered Delegate[®] WG product label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Refer to product label for Delegate for General Use Precautions, Mixing Directions, and Application Directions. Follow all use precautions and restrictions specified on the label.

Directions for Suppression of Spotted Wing Drosophila in Blueberries

Application Timing: Treat when pests appears and crop is becoming susceptible to spotted wing drosophila (SWD). Fruits become more susceptible as harvest approaches. Later harvested cultivars are at greater risk from SWD. Heavy infestations may require repeat applications but follow resistance management guidelines. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply 3 to 6 oz of Delegate WG per acre as a foliar spray to control spotted wing drosophila. The amount of Delegate WG applied per acre will depend upon plant size and volume of foliage present and pest pressure. Use a lower rate in the rate range for light infestations and/or small plants and a higher rate in the rate range for heavy infestations and/or larger plants. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Do not make more than three applications of Group 5 insecticides for thrips in a season. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Special Use Restrictions for Preharvest Interval of one day:

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- Do not apply more than a total of 17.9 oz of Delegate WG (0.281 lb ai spinetoram) per acre per year.
- **Maximum Number of Applications:** In order to utilize a 1-day Preharvest Interval, no more than three applications can be made per calendar year. See Resistance Management regarding number of applications for specific pests. See Resistance Management and Special Use Restrictions regarding number of applications for specific pests and locations.
- **Minimum Treatment Interval:** Do not make application less than 6 days apart between the first and second applications. Do not make applications less than 12 days apart between the second applications and the third application. The third application is applied at 1 day before harvest.

24(c) Registrant:
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268
1- 800-992-5994 (option #4)

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R359-029

Accepted: 08/15/17

Replaces: Initial printing