

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Delegate™ WG

EPA Reg. No. 62719-541

Foliar Insect Suppression in Dates

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Delegate™ WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Delegate WG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate WG.

Directions for Use

Refer to product label for General Use Precautions, Mixing, and Application information.

Pests and Application Rates:

Pest	Delegate WG (oz/acre)
carob moth	7

Application Timing: Apply Delegate WG when pests appear or in accordance with local conditions. Applications should closely follow regional spray recommendations. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional use recommendations for your area.

Resistance Management: Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- Do not apply more than a total of 19.5 oz of Delegate WG (0.305 lb ai spinetoram) per acre per year.
- **Maximum Number of Applications:** Do not make more than 4 applications per acre per calendar year.
- **Minimum Re-treatment Interval:** Do not make applications less than 7 days apart.
- **Preharvest Interval:** Do not apply within 7 days of harvest.

Note: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

R359-010
EPA accepted 08/13/09
Initial printing.

™ Trademark of Dow AgroSciences LLC