

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Delegate[®] WG EPA Reg. No. 62719-541

24(c) Special Local Need Registration SLN FL-090009

Low Volume Application in Citrus (For Distribution and Use Only in Florida)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read this SLN labeling and the label affixed to the container for Delegate[®] WG insecticide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the product container must be followed

Active Ingredient:
spinetoram (a mixture of
spinetoram-J and spinetoram-L)25.0%
Other Ingredients75.0%
Total100.0%

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Citrus

Citrus Fruits (Crop Group 10), including: Grapefruit, Lemons, Limes, Oranges, Tangerines

Pests and Application Rates:

(numbers in parentheses (-) refer to footnotes below)

Pets	Delegate WG (oz/acre)
Asian citrus psyllid(1)	3 - 6

(1) Control may be improved by the addition of an adjuvant to the spray mixture.

Delegate WG may be applied by ground in a minimum finished spray carrier volume of 2 gallons per acre, at rates recommended on the product label. Spray applications must be made with spray nozzles that create a droplet size with a volume median diameter (VMD) of 90 microns or larger. An adjuvant may be used in the spray mixture at the recommended rate of the manufacturer. The effect of specific adjuvants or tank-mix partners on Delegate WG performance in low volume applications should be evaluated in a small-scale application before making large-scale applications. See the product label on the Delegate WG container for additional information on tank-mixing and adjuvants.

To achieve good control, thorough coverage and proper equipment calibration are critical when using low spray volumes. Applications should only be made with air-assisted equipment. Low volume applications may not provide the same level of efficacy that can be obtained with an airblast sprayer at higher application volumes.

Reduce the potential for drift when making low volume applications. Factors that may impact drift include but are not limited to, wind speed and temperature inversion. Only apply this product if the wind direction favors on-target deposition. Do not apply when wind velocity exceeds 15 mph. See the product label on the Delegate WG container for additional information on reducing spray drift.

Restrictions:

- Do not apply more than a total of 12 oz of Delegate WG (0.188 lb ai of spinetoram) per acre per year.
- **Maximum Number of Applications:** Do not make more than 3 applications per calendar year.
- **Minimum Re-Treatment Interval:** Do not make applications less than 7 days apart.
- **Preharvest Interval (PHI):** Do not apply within 1 day of harvest.

®Trademark of Dow AgroSciences LLC
R359-015
Accepted: 07/14/09
Replaces R359-012