

Dimetric[®] EXT

Herbicide

EPA Reg. No. 1381-197
EPA SLN No. SD-170002

For Distribution and Use Only in the State of South Dakota

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, cancelled or suspended.

Read and understand the Special Conditions and Risks of Use Section on this label before using this product in accordance with this FIFRA Section 24(c) Special Local need.

Dimetric[®] EXT for use as a preemergence application for the suppression and control of certain broadleaf weeds in lentils and peas.

ATTENTION

- It is a violation of the Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN and the label affixed to the container for Dimetric[®] EXT herbicide before applying.

Use of Dimetric[®] EXT herbicide according to this SLN is subject to all use directions, precautions and limitations imposed by the label affixed to the container for Dimetric[®] EXT.

LENTILS AND PEAS

PREEMERGENCE APPLICATION: Make a single preemergence application of Dimetric EXT at 1/4 to 1/3 lb per acre per crop year. Apply in 10 or more gallons of water per acre with ground spray equipment. Apply Dimetric EXT before planting. Thorough incorporation, either by rainfall or by mechanical means, is essential for weed suppression. Under dry conditions, incorporate Dimetric EXT into the top 1 to 2 inches of soil with spike harrows, or similar shallow incorporation equipment, then cross harrow to ensure uniform soil incorporation. Where soil surface is moist at the time of application and rain follows before weed emergence, a broadcast application should provide adequate weed suppression.

Use the higher rate on fine-textured soils (high in clay or organic matter) and in fields with a history of high weed populations.

SPECIAL PRECAUTIONS: Crop injury may result if crop is under stress conditions caused by cold weather, low fertility, disease or insect damage. Crop injury may also result if application is followed by heavy rain.

Do not use on coarse-textured soils, sandy soils or soils with less than 1.5% organic matter.

Do not apply to "Estin" lentils.

Do not use on clay knobs or poorly covered subsoils.

Do not apply on shallow seedings less than 2 inches deep (preemergence only).

Maintain continuous spray tank agitation to keep material in suspension. Avoid overlapping and shut off spray booms while turning, slowing or stopping, or crop injury will occur.

NOTE: This treatment may cause some chlorosis or minor necrosis. Because lentil and pea varieties may vary in their susceptibility to Dimetric EXT, determining crop tolerance prior to adoption as a field scale practice is suggested to prevent possible injury.

Distributed by:

Winfield Solutions, LLC

PO 64589

St. Paul, MN 55164-0589

1/0330/7R₂ (SD-SLN)