

TECHNICAL INFORMATION BULLETIN

Restricted Use Pesticide. Due to toxicity to aquatic invertebrate animals. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP 15 INSECTICIDE



EPA Reg. No. 400-461

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency

FOR USE AND DISTRIBUTION IN THE STATE OF CALIFORNIA ONLY.

The addition of YELLOWSTRIPED ARMYWORM control in RICE.

Insect Growth Regulator

Not for Homeowner/Residential Use

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **DIMILIN® 2L Insecticide (EPA Reg. No. 400-461)**.

062917V071CA2ee

Crop	Pest	Application Rate (fl oz/acre)	Application Timing
RICE	<p>RICE RESTRICTIONS: Do not apply more than 16 fl oz of DIMILIN 2L (4 oz diflubenzuron ai) per acre per calendar year.</p> <p>Pre-harvest Interval: Do not harvest within 80 days of application. Do not use on rice fields in which crayfish (crawfish) farming is included in the cultural practice. Do not drain treated water into fields where crayfish farming is intended. Do not apply to rice immediately adjacent to sites of crayfish aquaculture. Do not use treated rice flood waters for irrigated crops except for uses currently established for DIMILIN 2L. Do not impregnate on granular materials. Do not use on wild rice (<i>Zizania</i> spp.).</p>		
	Yellowstriped Armyworm	4 - 8	Apply DIMILIN 2L at 4 - 8 fluid ounces per acre to rice when Yellowstriped armyworm eggs or larvae are first observed on plants. Use 4 fluid ounces per acre for small larvae (first to third instar), and 8 fluid ounces per acre for large larvae (fourth to sixth instar).
	<p>Consult your local extension service or UC IPM online for determination of economic threshold. DIMILIN 2L will prevent Yellowstriped armyworm larvae from successfully molting. DIMILIN 2L does not control adult Yellowstriped armyworm moths. DIMILIN 2L may have a transovarial effect on adult females causing them to produce unviable eggs.</p> <p>Aerial application: Apply DIMILIN 2L by air using at least 5 gallons total volume per acre.</p> <p>Ground application: Apply DIMILIN 2L by ground using at least 15 gallons total volume per acre.</p> <p>Do not apply DIMILIN 2L if flooding is in progress. Activity will be reduced. Since DIMILIN 2L is water active, the entire field must be treated.</p> <p>For maximum activity of DIMILIN 2L, do not disturb flood after a single application for at least 7 days. In water seeded, pinpoint or continuous flood rice, flood must not be disturbed for a minimum of 4 days following the first treatment. Hold treated water at least 14 days to allow for dissipation of DIMILIN 2L.</p> <p>DIMILIN 2L is not phytotoxic to rice. DIMILIN 2L can be safely applied in combination with post-permanent flood herbicides such as FACET®, GRANDSTAND® and LONDAX®. However, before using a tank-mix combination, read each product label carefully and follow Precautionary Statements on each label.</p>		

Manufactured For:
MacDermid Agricultural Solutions, Inc.
c/o ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

DIMILIN is a registered trademark of an Arysta LifeScience Group Company.

FACET is a registered trademark of BASF.

GRANDSTAND is a registered trademark of Dow AgroSciences LLC.

LONDAX is a registered trademark of United Phosphorus, Inc.

©2017 Arysta LifeScience Group Company.