



# Domark<sup>®</sup>

## 230 ME Fungicide

**Active Ingredient:**

Tetraconazole* .....	20.5%
<b>Other Ingredients:</b> .....	79.5%
<b>TOTAL:</b> .....	100.0%

\*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)propyl]1H-1,2,4-triazole

FOR CONTROL AND/OR SUPPRESSION OF CERTAIN DISEASES IN SOYBEAN & CORN

Domark<sup>®</sup> 230 ME Fungicide is a micro emulsion containing 1.9 pounds of tetraconazole per gallon.

EPA Registration No. 80289-7  
EPA Establishment No. 5905-IA-01

Batch Code will be placed on the container.

### KEEP OUT OF REACH OF CHILDREN CAUTION

**SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>If inhaled:</b>	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
<b>If in eyes:</b>	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
<b>If on skin or clothing:</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

(continued)

**FIRST AID (continued)**  
**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have affected person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

For Health Emergency or Spill (24 hr) call 800-892-0099 or for Transportation Chemical Emergency Spill Leak Fire Exposure or Accident call CHEMTREC Day or Night 800-424-9300 or 202-483-7616

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, and viton ≥ 14 mils. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any water proof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENVIRONMENTAL HAZARDS:

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise caution when making applications of *Domark*, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment wash waters or rinsate.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Do not** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all activities with the exception of 20 days for detasseling corn grown for seed. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any water proof material
- Shoes plus socks

### PRODUCT INFORMATION

*Domark* fungicide is formulated as a 1.9 pound active ingredient per gallon micro emulsion (ME). The active ingredient in *Domark* is tetraconazole, a triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. Tetraconazole is absorbed quickly into the plant tissue and like all triazoles can move up, but not down the plant. Optimal disease control is achieved when *Domark* is applied in a regularly scheduled spray program. When using *Domark* in combination and/or rotation with other fungicides, it is

important to use fungicides that have different modes of action (i.e. non Group 3 fungicides). Since *Domark* is a sterol biosynthesis inhibiting fungicide, do not rotate with other sterol biosynthesis inhibitors, such as Folicur<sup>®</sup>, Tilt<sup>®</sup> or Laredo<sup>®</sup>

### MODE OF ACTION

The active ingredient in *Domark* is tetraconazole, which belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the Fungicide Resistance Action Committee (FRAC) as Target Site of Action Group 3 fungicides.

### RESISTANCE MANAGEMENT

*Domark* contains tetraconazole, a Group 3 fungicide (sterol biosynthesis inhibitors), and is effective against labeled pathogens resistant to fungicides with modes of action different from those of target site Group 3, such as dicarboximides, strobilurins, benzimidazoles, or phenylamides. However, fungal isolates resistant to Group 3 fungicides may eventually dominate the fungal population if Group 3 fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species, especially if resistance to Group 3 fungicides is already present in the pathogen population. This may result in reduced disease control by Group 3 fungicides. To maintain the performance of *Domark* in the field, do not exceed the total number of sequential applications of *Domark* and the total number of applications of *Domark* per year stated in "CROP USE RATES AND TIMING OF APPLICATIONS". Adhere to the label instructions regarding the consecutive use of *Domark* or other target site of action Group 3 fungicides that have a similar site of action on the same pathogens. Consider the following to delay the development of fungicide resistance:

1. **Tank mixtures:** If *Domark* is used in tank mixtures with fungicides from different mode of action Groups that are registered for the same use and that are effective against the pathogens of concern, use at least the minimum labeled rates of each fungicide in the tank mix.
2. **IPM:** Integrate *Domark* into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or Valent representative for additional IPM strategies established for your area. Use *Domark* in Agricultural Extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
4. **Reporting:** If a Group 3 target site fungicide appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact your Valent representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

## RAINFASTNESS

*Domark* is rainfast 2 hours after application. **Do not** apply if rain is expected within 2 hours of application or disease control may be reduced.

## JAR TEST TO DETERMINE COMPATIBILITY OF *DOMARK*

Perform a jar test before mixing commercial quantities of *Domark* when using *Domark* for the first time, or when a new water source is being used.

1. Add 1 pt of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 ml of *Domark* to the quart jar; gently mix until product goes into suspension.
3. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
4. An ideal tank mix combination will be uniform and free of suspended particles.

## SPRAYER PREPARATION

Before applying *Domark*, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *Domark*. If two or more products were tank mixed prior to *Domark* application, follow the most restrictive cleanup procedure.

## SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, observe them. Do not apply this product when weather conditions favor spray drift from treated areas. When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION".

## MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the *Domark* to the spray tank. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing *Domark* with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions.
4. Fill spray tank to desired level with water. Agitation should continue until all spray solution has been applied.
5. Mix only the amount of spray solution that can be applied the day of mixing. *Domark* should be applied within 24 hours of mixing.
6. When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. No label dosage rates should be

exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

7. Do not combine *Domark* in a sprayer tank with pesticides or fertilizers, unless your prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use.

## APPLICATION EQUIPMENT

Application equipment must be clean and in good condition. Frequently check nozzles for accuracy.

## SPRAYER CLEANUP

Clean spray equipment each day following *Domark* application. After *Domark* is applied, use the following steps to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

Thoroughly clean spray equipment, including all tanks, hoses, booms, screens and nozzles, before it is used to apply foliar pesticides.

## AERIAL APPLICATION

To avoid drift, apply the largest droplet size possible that will provide uniform coverage and result in satisfactory disease control. To obtain satisfactory application and avoid drift, the following directions must be observed:

**Do not** apply during low-level inversion conditions, when winds are gusty or under other conditions that favor drift. Application should be avoided when wind velocity is less than 2 mph and more than 15 mph.

### • Carrier Volume and Spray Pressure:

- For aerial application use a minimum of 2 gallons per acre for all diseases except rust and white mold/*Sclerotinia* stem rot of soybeans for which a minimum of 5 gallons per acre must be used. Increasing the spray volume to 7 gallons or more per acre generally provides better coverage and more consistent disease control.

- **Do not** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- **Nozzle Selection and Orientation:** Minimize formation of very small drops by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles, such as diaphragm type nozzles, to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. **Do not** place nozzles on the outer 25% of the wings or rotors.



### **GENERAL CHEMIGATION INSTRUCTIONS:**

Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other irrigation experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### **Requirements for Chemigation Systems Connected to Public Water Systems**

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventor (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm

pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add *Domark* slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

*Domark* should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

### **Sprinkler Chemigation:**

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add *Domark* slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

*Domark* should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

### ROTATIONAL CROP RESTRICTIONS

There are no time interval restrictions for the tetraconazole use approved crops peanut, pecan, soybean and sugar beet. For other crops use the time intervals listed on right to determine the minimum required time interval between the last *Domark* application and new crop planting.

Rotational Crop Guideline	
Crop	Time Interval in Days Before Planting
Small Grains (barley, buckwheat, millet, oats, rice, rye, triticale and wheat)	45
Sugarcane	45
All Other Crops	120

### CROP USE RATES AND TIMING OF APPLICATIONS

Field Corn, Popcorn, Corn Grown For Seed Production				
Disease	Dosage Rate		When to Apply	Special Use Instructions
	Fl oz/A	GPA		
Gray Leaf Spot ( <i>Cercospora zeae-maydis</i> )	4.0 to 6.0 (0.060 to 0.090 lb ai/A)	Ground: Minimum of 10 GPA  Aerial: Minimum of 2 GPA	Apply prior to disease onset when conditions favor disease development. Curative applications are most effective when disease incidence does not exceed 5% of the plants at time of application.	Use <i>Domark</i> as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants. To limit the potential for resistance development, do not apply more than 6 fl oz per acre per year.
Rust, Common ( <i>Puccinia sorghi</i> )				
Rust, Southern ( <i>Puccinia polysora</i> )				
Anthraxnose Leaf Blight ( <i>Colletotrichum graminicola</i> )				
Eye Spot ( <i>Aureobasidium zeae</i> )				
Northern Corn Leaf Blight ( <i>Exserohilum turcicum</i> )				
Northern Corn Leaf Spot ( <i>Bipolaris zeicola</i> )				
Southern Corn Leaf Blight ( <i>Bipolaris maydis</i> )				

### RESTRICTIONS AND LIMITATIONS

1. **Do not** make more than (1) application per year.
2. **Do not** apply more than 6 fl oz (0.090 lb ai tetraconazole) of *Domark* per acre per year.
3. **Do not** apply *Domark* after corn growth stage R-3 (milk).
4. **Do not** use adjuvants in sprays made between V-8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. A compatibility agent, another fungicide, or an insecticide may be included if needed and labeled for use in corn. Refer to adjuvant product label for specific use directions and restrictions. Always follow the more restrictive label.

## CROP USE RATES AND TIMING OF APPLICATIONS

<b>Soybean</b>				
<b>Disease</b>	<b>Dosage Rate</b>		<b>When to Apply</b>	<b>Special Use Instructions</b>
	<b>Fl oz/A</b>	<b>GPA</b>		
Asian Soybean Rust ( <i>Phakopsora pachyrhizi</i> )	4.0 to 5.0 (0.060)	Ground: Minimum of 10 GPA	Apply prior to disease development when infections are likely to occur.	Use <i>Domark</i> as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of soybeans.
Cercospora Blight ( <i>Cercospora kikuchii</i> )	to 0.075 lb ai/A)	Aerial: Minimum of 2 GPA;	If necessary repeat with a second application before growth stage R-6.	
Purple Seed Stain ( <i>Cercospora kikuchii</i> )		(5 GPA for White Mold and Asian Soybean Rust)	Curative applications are most effective when disease incidence does not exceed 5% of the soybean plants at time of application.	
Frogeye Leaf Spot ( <i>Cercospora sojina</i> )			Make application at soybean growth stage R-3 (early pod fill) or when conditions are favorable for disease development. Repeat application 15 to 21 days after first application if disease pressure is heavy.	
White Mold/Sclerotinia Stem Rot ( <i>Sclerotinia sclerotiorum</i> )			Under severe disease conditions the higher rate and shorter spray intervals should be used.	
Powdery Mildew ( <i>Microsphaera diffusa</i> )				
Brown Spot ( <i>Septoria glycines</i> )				
Anthracnose ( <i>Colletotrichum</i> spp.)				

### RESTRICTIONS AND LIMITATIONS

1. **Do not** make more than two (2) applications per year.
2. **Do not** apply more than 10 fl oz (0.150 lb ai tetraconazole) of *Domark* per acre per year.
3. **Do not** graze or feed *Domark*-treated forage or hay to livestock.
4. **Do not** apply *Domark* after soybean growth stage R-5 (beginning seed).
5. **Do not** harvest immature soybeans for consumption once plants are treated with *Domark*.
6. **Do not** use on vegetable soybean varieties grown for their immature pods.

*Domark* may be tank mixed with the following products for use in soybeans:

#### **Herbicides:**

glyphosate (i.e. Roundup®)  
clethodim (i.e. Select Max®)

#### **Fungicides:**

azoxystrobin (i.e. Quadris®)  
pyraclostrobin (i.e. Headline®)

#### **Insecticides:**

acephate (i.e. Orthene®)  
cyfluthrin (i.e. Baythroid®)  
gamma-cyhalothrin (i.e. Proaxis®)  
permethrin (i.e. Pounce®)

#### **Insecticides (continued):**

chlorpyrifos (i.e. Lorsban®)  
esfenvalerate (i.e. Asana®)  
lambda-cyhalothrin (i.e. Warrior®)  
zeta-cypermethrin (i.e. Mustang Max™)

#### **Use Restrictions for *Domark* Tank Mixes:**

1. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.
2. The most restrictive labeling applies when using a tank mix.

## WARRANTY STATEMENT

Read the entire label before using this product, including this Limitation of Warranty and Liability.

If the terms are not acceptable, return the product at once unopened for a refund of the purchase price.

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks described below, when used in accordance with the Directions for Use under normal conditions.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, VALENT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Buyers and Users of this product must be aware that there are inherent unintended risks associated with the use of this product, independent from the control of Valent. These risks include, but are not limited to, weather conditions, soil factors, moisture conditions, irrigation practices, condition of the crop at the time of application, materials which are present in the tank mix with this product or prior to the application of it, cultural practices or the manner of use or application, all risks which are impossible to eliminate. The Buyers and Users must be aware that these factors may cause: ineffectiveness of the product, reduction of harvested yield of the crop (entirely or partially), or crop injury.

If the Buyer does not agree with the acceptance of these risks, then **THE PRODUCT SHOULD NOT BE APPLIED.** To the extent consistent with applicable law, the Buyer acknowledges and accepts these inherent unintended risks and **AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.**

To the extent consistent with applicable law, Valent or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise). To the extent consistent with applicable law, the exclusive remedy of the User or Buyer and the exclusive Liability of Valent or Seller shall be the return of the purchase price of the product, or at the election of Valent or Seller, the replacement of the product.

To the extent that applicable law allows such requirements, Valent or its Seller must have prompt notice of any claim so that an immediate inspection of Buyer's or User's growing crops can be made. To the extent consistent with the applicable law, if Buyer and User do not notify Valent or Seller of any claims, in proper time, it shall be barred from obtaining any remedy.

To the extent consistent with applicable law, Buyers and Users are deemed to have accepted the terms of this Limitation of Warranty and Liability, which may not be modified by any verbal or written agreement.

---

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

### PESTICIDE STORAGE

Store in original container in a dry, temperature-controlled, secure, place.

### PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

**For rigid, non-refillable containers (2.5 to 5 gallons):** Nonrefillable container. Do not reuse or refill the container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of resulting smoke.

ESL 083011

REV 100511

*Domark* is a registered trademark of Isagro S.p.A.  
*Select Max* is a registered trademark of Valent U.S.A. Corporation  
Asana is a registered trademark of E.I. duPont de Nemours and Company  
Baythroid is a registered trademark of Bayer Crop-Science  
Folicur is a registered trademark of Bayer Crop-Science  
Headline is a registered trademark of BASF  
Laredo is a registered trademark of Dow AgroSciences, LLC  
Lorsban is a registered trademark of Dow AgroSciences, LLC  
Mustang Max is a trademark of FMC Corporation  
Orthene is a registered trademark of OMS Investments, Inc.  
Pounce is a registered trademark of FMC Corporation  
Proaxis is a trademark of Pytech Chemicals GmbH  
Quadris is a registered trademark of Syngenta  
Roundup is a registered trademark of Monsanto Company  
Tilt is a registered trademark of Syngenta  
Warrior is a registered trademark of Syngenta

Manufactured by Isagro S.p.A. for:



**Isagro USA, Inc.**  
430 Davis Drive, Suite 240  
Morrisville NC 27560  
Made in U.S.A.  
Form 1562-F  
EPA Registration No. 80289-7  
EPA Establishment No. 5905-IA-01

Distributed by:  
**Valent U.S.A. Corporation**  
P.O. Box 8025  
Walnut Creek CA 94596

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-6-VALENT (682-5368).



For state registration and/or supplemental labels, please call or visit us online.

*Products That Work, From People Who Care*® | [www.valent.com](http://www.valent.com) | 800-6-VALENT (682-5368)

***Always read and follow label instructions.***

©2011 Valent U.S.A. Corporation. All rights reserved. Printed in the U.S.A.