

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### **DUPONT™ CORAGEN® INSECT CONTROL**

EPA Reg. No. 352-729

### **FOR CONTROL OF COLORADO POTATO BEETLE AND CABBAGE LOOPERS IN POTATOES IN THE STATES OF IDAHO, NEVADA, OREGON, AND WASHINGTON**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ CORAGEN® is recommended for the control of Colorado Potato Beetle and Cabbage Looper larvae in potatoes.

Apply CORAGEN® at 2.0 to 5.0 fluid ounces per acre as a foliar spray or via overhead chemigation. For best results with foliar sprays, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v). For chemigation applications, apply in 0.1 to 0.2 acre inches of water and add MSO at 12 to 16 fl oz/acre.

Apply when populations reach local established thresholds to prevent crop damage.

#### **CABBAGE LOOPER**

CORAGEN® applied at 2.0 to 3.0 fluid ounce per acre rate should be used on early stage instars of Cabbage Looper (1st through 3rd instar). Larger larvae may require higher rates.

#### **COLORADO POTATO BEETLE**

CORAGEN® applied at 2.0 to 3.0 fluid ounces per acre should be used where local populations of Colorado Potato Beetle are believed to be sensitive to most commonly used insecticides. Apply just prior to or just after egg hatch while larvae are small. In some areas, where local populations of Colorado Potato Beetle have elevated levels of resistance to insecticides, CORAGEN® should be used at the 5.0 fluid ounce rate. With resistant populations of Colorado Potato Beetle, back-to-back applications on 5 to 7 day intervals may be required to achieve maximum control.

Do not make more than two sequential applications of CORAGEN® before rotating to another effective registered insecticide having a different mode-of-action.

See the EPA-registered label for complete instructions on overhead chemigation.

#### **USE RESTRICTIONS**

Do not apply more than 15.4 fluid ounces of CORAGEN® per acre per year.

Do not make more than 4 applications per acre per crop.

The minimum retreatment interval is 5 days.

The preharvest interval is 14 days.

The field re-entry interval is 4 hours.

Read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

(Replaces R-1821 011217 01-16-17)

**IMPORTANT NOTICE**

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A