

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### DUPONT™ CORAGEN® INSECT CONTROL

EPA Reg. No. 352-729

### TRANSPLANT WATER TREATMENT FOR SUPPRESSION OF CABBAGE MAGGOT IN CABBAGE IN THE STATE OF NEW YORK

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ CORAGEN® insect control is recommended for the suppression of Cabbage Maggot in Cabbage.

Apply DuPont™ CORAGEN® insect control at 5.0 fluid ounces per acre as a transplant water treatment at planting using properly calibrated application equipment for control of Cabbage Maggot (*Delia radicum*).

**SOIL APPLICATIONS** (transplant water treatment): CORAGEN® must be applied uniformly in the root zone or poor performance will result.

#### Transplant water treatment:

Transplants should be adequately watered before transplanting. Apply at transplanting in a minimum of 2 fluid ounces of treatment solution per transplant. Ensure water volume is sufficient to thoroughly wet the root zone.

#### Important

Before using DuPont™ CORAGEN® insect control read and follow all applicable directions, restrictions and precautions on the EPA-registered label.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

#### Precautions

Make no more than 4 applications per acre per crop.

Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per crop.

Do not apply more than 10 fl oz (0.132 lb ai per acre) of CORAGEN® per crop by any combination of at plant soil application and drip chemigation.

Preharvest interval (PHI): Last application (days to harvest) is 3.

The field re-entry interval (REI) is 4 hours.

(Replaces R-1533 121014 01-08-15)

#### IMPORTANT NOTICE

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A