

# FIFRA Section 24(c) Special Local Need (SLN) Label

DuPont Crop Protection

Chestnut Run Plaza, 974 Centre Road

Wilmington, DE 19805 USA

## Curzate® 60DF

EPA Reg. No. 352-592

EPA Section 24(c) Special Local Need Registration SLN No. OR-150004

### For Control of Downy Mildew in Spinach, Cabbage, Swiss Chard and Table Beet Grown for Seed

This label is valid until December 31, 2024 or until otherwise amended, withdrawn, canceled, or suspended

(For Distribution and Use Only in the State of Oregon)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Curzate® 60DF before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as described in this supplemental labeling, use of Curzate 60DF is subject to all precautions and limitations imposed by the label affixed to the product container.

Active ingredients	By Weight
Cymoxanil.....	60%
Other ingredients .....	40%
<b>TOTAL</b> .....	<b>100%</b>

#### KEEP OUT OF REACH OF CHILDREN

#### WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

#### DIRECTIONS FOR USE

Curzate® 60DF fungicide may be applied to spinach, cabbage, Swiss chard and table beet grown for seed production only for the control of downy mildew. Consult the Curzate 60DF EPA-registered label for additional requirements, restrictions and precautions.

#### HOW TO USE

##### Use Rate, tank mixes and timing

- Apply Curzate 60DF fungicide at 3.2 ounces per application per acre in sufficient water to obtain thorough spray coverage of the crop.
- Use Curzate 60DF only in combination with another registered protectant fungicide, such as mancozeb or chlorothalonil.
- Applications can be made prior to or after transplanting.
- Initial applications should start when local conditions indicate that downy mildew infection is imminent.
- Make applications on a 5-7 day interval as long as disease conditions exist.
- Do not exceed 5 applications of Curzate 60DF per 12 month period.

- For conventional ground application equipment, including tractor-mounted sprayers, boom or backpack sprayers, use a minimum of 20 gals/acre.
- For hard-to-wet foliage, a spray adjuvant may be used to help increase spray coverage and retention.

**Restrictions/Precautions:**

- Do not apply by air.
- For use under this SLN label do not apply this product through any type of irrigation system.
- For use only on specified crops grown for seed.
- Do not replant treated fields to any crop within 30 days after- last Curzate 60 DF treatment.

**Special Crop Use Restrictions**

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057- 0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies.”

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

R612-002

Accepted: 01/09/2020

Replaces R-1534 060415-2