

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

FOR CONTROL OF KATYDID* NYMPHS IN CITRUS IN THE STATE OF CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Exirel® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont EXIREL® insect control, is recommended for control of katydid nymphs in citrus.

Apply EXIREL® insect control at 16.0 to 20.5 fluid ounces/acre plus adjuvant (see adjuvant section of the Federal Label for more details) as a foliar spray using properly calibrated ground spray equipment. Apply at first indication of katydid nymphs. Initial application should be made at first sign of infestation to the early nymphal stage before pest reaches damaging levels. Monitor fields and make additional application if populations begin to rebuild.

Thorough coverage of leaves and fruit is essential to achieve best results. Use the higher application rate for moderate to heavy insect pressure. Where higher spray volumes are used (greater than 150 gallons per acre), apply a higher EXIREL® rate in the specified rate range. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for this pest.

Read and follow the protection of pollinators section of the main label when using during bloom.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

RESTRICTIONS

Do not make apply more than 0.4 lbs ai/A of CYAZYPYR® or cyantraniliprole containing products per year.

The minimum retreatment interval is 7 days

The pre-harvest interval for citrus is 1 day

The field re-entry interval for citrus is 12 hours

*Forktailed bush katydid (*Scudderia furcata*)

(Replaces R-1713 022516 02-25-16)

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

R-4175 050919 12-11-18

Page 1 of 1

Expires 5/31/21