



2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

FOR THE CONTROL OF CITRUS THRIPS IN HIGH AND LOWBUSH BLUEBERRIES IN THE STATE OF CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ EXIREL® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont™ Exirel® is recommended for control of citrus thrips in blueberries.

Apply DuPont™ Exirel® at 13.5 to 20.5 fluid ounces per acre as a foliar spray.

For best performance use an effective adjuvant. Read the “Use of Adjuvants” and “Tank Mixtures and Crop Safety” sections of the main EXIREL® label.

Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of bushes or plants and density of foliage.

For best results, apply 100-150 gallons of water per acre when using commercial airblast equipment. Do not apply less than 30 gallons of water per acre when using commercial airblast equipment.

Use Exirel® as part of an effective thrips control program. Rotate with products with different modes of action. Begin making applications early in the pest development cycle when thrips populations are low. If populations are above local established threshold, use an effective thrips knockdown product before applying DuPont™ Exirel®.

Read and follow the protection of pollinators section of the main EXIREL® label when using during bloom.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

RESTRICTIONS

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® or cyantranilprole containing products per acre per calendar year.

Re-entry interval is 12 hours.

Pre-harvest interval is 3 days.

(Replaces: R-1634 070915 07-09-15)

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A