

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ EXIREL® INSECT CONTROL

EPA Reg. No. 352-859

FOR CONTROL OF APPLE LEAFCURLING MIDGE IN APPLES IN THE STATES OF CALIFORNIA, MAINE, MASSACHUSETTS, OREGON, NEW YORK, AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Exirel® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont™ EXIREL® Insect Control is recommended for the control of Apple Leafcurling Midge (*Dasineura mali*).

Apply EXIREL® at 10 to 20 fluid ounces per acre at first sign of egg laying or at early egg hatch. Two sequential applications, 10 to 14 days apart, may be needed in areas of high pest pressure.

Application should be made as a thorough coverage foliar spray using standard airblast ground equipment. The addition of a high quality spreader/penetrator adjuvant (such as horticultural mineral oil, see adjuvant section of the Federal Label for more details) may improve coverage and penetration into leaf tissue.

Read and follow the protection of pollinators section of the main label when using during bloom.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

RESTRICTIONS

Do not make apply more than 0.4 lbs ai/A of CYAZYPYR® or cyantraniliprole containing products per year.

Make no more than 3 applications of EXIREL® or other Group 28 insecticides within a single generation of the target pest on a crop.

The minimum retreatment interval is 7 days

The pre-harvest interval for apples is 3 days

The field re-entry interval for apples is 12 hour

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A