

RESTRICTED USE PESTICIDE DUE TO HIGH ACUTE TOXICITY TO HUMANS

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application; mixing; loading or repair or cleaning of application equipment.

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ LANNATE® SP INSECTICIDE

EPA Reg. No. 352-342

FOR CONTROL OF SPOTTED ALFALFA APHID ON ALFALFA IN THE STATES OF OKLAHOMA, NEW MEXICO, KANSAS, COLORADO, NEBRASKA, WYOMING AND CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

DuPont™ LANNATE® SP insecticide is recommended for the control of spotted alfalfa aphid in alfalfa. Apply LANNATE® SP at 0.5 to 1 pounds/acre when the local threshold is reached.

APPLICATION

Apply by properly calibrated air or ground equipment using sufficient spray volume to obtain thorough coverage. Make repeat applications as needed based on scouting or on a 5 to 7 day schedule.

Chemigation: LANNATE® SP may be applied by overhead sprinkler chemigation. For best results use the highest listed rate of LANNATE® SP. Apply in 0.1 to 0.2 inches of water per acre. See chemigation section of main label for more information.

NOTE:

Do not apply to dormant or semidormant alfalfa when minimum daily temperature is 50 oF or lower.

Do not apply more than 4 pounds of LANNATE® SP per acre per crop.

Do not make more than 10 applications per crop.

The field re-entry interval following application of LANNATE® SP is 48 hours.

Do not apply within 7 days of cutting or allowing livestock to graze.

When LANNATE® SP is used on alfalfa grown for seed, the seed may not be used for sprouts. All seed from treated crop must be tagged, “Not for Human Use” at the processing plant.

(Replaces R-1382 071813 07-19-13)

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A