

**RESTRICTED USE PESTICIDE**

**DUE TO HIGH ACUTE TOXICITY TO HUMANS**

**For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading or repair or cleaning of application equipment.**



**2(ee) Recommendation**

Use information in accordance with Section 2(ee) of FIFRA

**DUPONT™ LANNATE® SP INSECTICIDE**

EPA Reg. No. 352-342

**FOR USE ON SWEET POTATOES FOR THE CONTROL OF  
WESTERN YELLOW-STRIPED ARMYWORM VIA GROUND  
FOLIAR SPRAY APPLICATIONS IN THE STATE OF  
CALIFORNIA**

**This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DuPont™ LANNATE® SP can be applied by ground application at 1 pound per acre when western yellow-striped armyworm populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

**APPLICATION**

LANNATE® SP should be applied at 5 to 7 day intervals to maintain control. Use sufficient water to obtain thorough, uniform coverage. Since LANNATE® SP is a fast acting contact insecticide, best results follow direct spraying of the target insect. Do not apply this product through any type of irrigation system.

**NOTE:**

Do not apply LANNATE® SP within 30 days of harvest.

Do not apply more than 2.7 pounds of methomyl (3 pounds of LANNATE® SP) per acre per growing season.

Do not make more than a total of 3 applications of LANNATE® SP per crop.

The reentry interval is 48 hours.

(Replaces R-1312 111612 12-18-12)

**IMPORTANT NOTICE**

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use. If the terms are not acceptable, return the product at once, unopened, for a refund of the purchase price. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT

or at <http://www.dupont.com/cgi-bin/ag/prodsearch/start.cgi>

**2016 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, Delaware 19805 U.S.A.**