



2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

DUPONT™ LEADOFF® HERBICIDE

EPA REG. NO. 352-853

PLUS DUPONT™ PANOFLEX™ HERBICIDE

EPA Reg. No. 352-876

OR ABUNDIT™ EXTRA HERBICIDE

EPA REG. NO. 71368-20

DUPONT™ LEADOFF® HERBICIDE REDUCED RATE FOR USE IN POST HARVEST OR FALLOW BURNDOWN OF WEEDS IN NORTH CAROLINA, SOUTH CAROLINA AND VIRGINIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

LEADOFF® Herbicide may be used at 1.0 ounce per acre and tank mixed with PANOFLEX™ Herbicide at 0.3 – 0.6 ounces per acre or ABUNDIT™ EXTRA Herbicide for burndown of weeds when applied post-harvest or to fallow fields. See PANOFLEX™ Herbicide and ABUNDIT™ EXTRA Herbicide labels for weeds controlled and weed size limitations. For improved weed control, LEADOFF® Herbicide may be tank mixed with other herbicides, such as 2,4-D, that are labeled for post-harvest or fallow fields.

DuPont will not be responsible for losses or damages resulting from the use of these products in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product.

SURFACTANTS

Include a spray adjuvant with applications. In addition, an ammonium nitrogen fertilizer may be used. For best results, add a crop oil concentrate or modified seed oil (methylated, ethylated or saponified) at 1% v/v (1 gallon per 100 gallons of spray) OR, add nonionic surfactant at 0.25 - 0.5% v/v (1 - 2 quarts per 100 gallons of spray).

ROTATIONAL CROP GUIDELINES

Corn may be planted immediately after a LEADOFF® Herbicide application or 14 days after a PANOFLEX™ Herbicide application.

Cotton or soybeans may be planted 30 days after application.

Peanuts or tobacco may be planted 45 days after application.

Refer to the appropriate product label for additional rotational crop guidelines

IMPORTANT PRECAUTIONS:

Refer to the LEADOFF®, PANOFLEX™, ABUNDIT™ EXTRA, and any tank mix partner labels for information on weed species controlled, additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on any label shall apply.

(Replaces: R-1507 010515 12-19-14)

© 2014 E. I. du Pont de Nemours and Company. All rights reserved.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2014 E. I. du Pont de Nemours and Company. All rights reserved.