

Supplemental Labeling



Dow AgroSciences

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

DuraCor™

EPA Reg. No. 62719-739

For Distribution and Use Only in the States of:

AL, AR, AZ, CO*, FL, GA, ID*, KS, KY, LA, MO, MS, MT*, ND*, NE, NV*, NM, OK, SD*, TN, TX, UT, WY*

**For Use on Grass Harvested for Hay
Intended for Distribution or Sale Off the Farm or Ranch**

**For Use on Grass Harvested for Silage, Haylage, Baylage, or Green Chop
Intended for Use On the Farm or Ranch**

* For use on rights-of-ways and utility lines transecting grazed areas.

This supplemental label expires on November 22, 2022, and must not be used or distributed after this date.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for DuraCor herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for DuraCor.

Use Precautions and Restrictions

<p>IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS</p> <ul style="list-style-type: none"> • Carefully read the section <i>Restrictions in Hay or Manure Use</i>. • It is mandatory to follow the Use Precautions and Use Restrictions on this label. • Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid and florpyrauxifen-benzyl to cause injury to sensitive broadleaf plants. • Hay can only be used on the farm or ranch where product is applied unless allowed by supplemental labeling. • The applicator must provide the land manager with a copy of the Dow AgroSciences stewardship instructions regarding uses of forage from areas treated with aminopyralid and florpyrauxifen-benzyl. • Consult with a Dow AgroSciences representative if you do not understand the Use Precautions and Use Restrictions. Call 800-258-3033 Customer Information Group. 	<p>Forage and Manure Management</p> <p>©Copyright 2011 Dow AgroSciences LLC</p>
---	--

- **Do not use grasses treated with DuraCor in the preceding 18 months for hay intended for export outside the United States.**
- **Do not use hay or straw from areas treated with DuraCor within the preceding 18 months or manure from animals feeding on hay treated with DuraCor in compost.**
- **Do not use grasses treated with DuraCor within the preceding 18 months for seed production.**
- **Grazing and Haying Restrictions:** Do not harvest forage for hay within 7 days of DuraCor application. Cutting hay too soon after spraying weeds can compromise the weed control. Wait 14 days prior to cutting grass hay to allow for maximum herbicide activity.
- **Transfer of Animals Feeding on DuraCor-Treated Forage:** Do not transfer animals grazing or feeding on hay from areas treated with DuraCor to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid and florpyrauxifen-benzyl to cause injury to sensitive broadleaf plants.
- **Seeding Legumes:** Do not plant forage legumes until a soil bioassay has been conducted to determine if aminopyralid and florpyrauxifen-benzyl residues remaining in the soil will adversely affect the legume establishment.
- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not allow livestock to graze treated areas until poisonous plants are dry and no longer palatable to livestock.

- **Restrictions in Hay or Manure Use:**
 - Do not use treated plant residues, including hay or straw from areas treated within the preceding 18 months, in compost, mulch or mushroom spawn.
 - Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch or mushroom spawn.
 - Do not spread manure from animals that have grazed or consumed forage or hay from treated areas within the previous 3 days on land used for growing broadleaf crops.
 - Manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated and florpyrauxifen-benzyl-treated areas within the previous 3 days may only be used on pasture grasses, grass grown for seed, wheat, and corn.
 - Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or eaten hay harvested from aminopyralid-treated and florpyrauxifen-benzyl-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid and florpyrauxifen-benzyl residue in the soil is at a level that is not injurious to the crop to be planted.
 - To promote herbicide decomposition, plant residues should be evenly incorporated in the surface soil or burned. Breakdown of aminopyralid and florpyrauxifen-benzyl in plant residues or manure is more rapid under warm, moist soil conditions and may be accelerated by supplemental irrigation.
- **Preharvest Interval:** Do not cut forage for hay within 7 days of application. For program lands, such as CRP, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Crop Rotation:** Do not rotate to cropland for one year following an application of DuraCor. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of aminopyralid and florpyrauxifen-benzyl present in the soil will not adversely affect that broadleaf crop.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern, or drainage. The field bioassay can be initiated one year after the last application of aminopyralid and florpyrauxifen-benzyl in that field. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, forage grasses, native grasses, or grasses grown for hay.
- **DuraCor is highly active against many broadleaf plant species.** Do not use this product on areas where loss of desirable broadleaf forage plants, including legumes, cannot be tolerated.
- Trees adjacent to or in a treated area can occasionally be affected by root uptake of DuraCor through movement into the soil. Do not apply DuraCor within the root zone of desirable trees unless such injury can be tolerated. Use special caution near roses and leguminous trees such as locusts, redbud, mimosa, and caragana.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R640-002

EPA Accepted: 11/26/19

Initial printing