

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Duramax<sup>®</sup>

EPA Reg. No. 62719-556

### Preharvest and Spot Treatments of Weeds in Dry Peas, Lentils, and Chickpeas (For Distribution and Use Only in the States of Idaho, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota and Washington)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

#### Preharvest Application

This product may be applied as a broadcast over the top spray to control labeled weeds prior to harvest of dry peas, lentils, or chickpeas. Apply up to 24 fl oz in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less).

#### Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- Make only one preharvest application per year. Do not combine a preharvest spray with a spot treatment on the same crop area.
- There is at least a 30-day plantback interval between treatment and replanting for any crop not listed in the label for this product.
- Do not feed treated vines and hay from these crops to livestock.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Do not apply preharvest to dry beans, peas, lentils, or chickpeas grown for seed as a reduction in germination or vigor may occur.
- Do not treat field (feed) peas or cowpeas since these are considered to be grown as livestock feed.

#### Spot Application

This product may be applied as a spot spray to control weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed. Apply up to 24 fl oz per acre in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, apply at or beyond bud/heading stage or growth of target weeds. **Note:** The crop receiving the spray in the spot treated area will be killed.

**Precautions and Restrictions:**

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- Make only one application per year. Do not combine a spot treatment with a preharvest spray on the same crop area.
- There is at least a 30-day plantback interval between treatment and replanting for any crop not listed in the label for this product.
- Do not feed treated vines and hay from these crops to livestock.
- Do not treat field (feed) peas or cowpeas since these are considered to be grown as livestock feed.

®™ Trademark of Dow AgroSciences LLC

R352-020

EPA accepted 11/26/10

Replaces R352-015