

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Durango<sup>®</sup> DMA<sup>®</sup>

EPA Reg. No. 62719-556

### Preharvest and Spot Treatments of Weeds in Dry Beans

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

Refer to the label booklet for this product for general information on product use, mixing, application equipment and techniques, labeled uses, and weed rate tables.

To minimize drift, use nozzles that emit medium to large-sized droplets. The likelihood of drift occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are meteorological conditions that favor spray drift. Extreme care must be used when applying this product to avoid injury to desirable plants and crops.

#### Broadcast Application

This product may be applied as a broadcast over-the-top spray to control labeled weeds prior to harvest of dry beans. Apply up to 24 3/4 fl oz in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Ground or aerial broadcast applications may be made.

#### Precautions and Restrictions:

- Apply at least 7 days before harvest.
- Make only one application per crop. Do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the label for this product.
- Do not apply preharvest for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay to livestock. Do not apply this product through any type of irrigation system.
- Do not treat cowpeas, since these are considered to be grown as livestock feed.

**Spot Application**

This product may be applied as a spot spray to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 24 3/4 fl oz in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a hand-held sprayer. For best results, applications should be made at or beyond bud/heading stage or growth of target weeds.

**Note:** The crop receiving the spray in the spot treated area will be killed.

**Precautions and Restrictions:**

- Apply at least 7 days before harvest.
- Make only one application per year. Do not combine a spot treatment with a preharvest spray on the same crop area.
- Employ at least a 30-day plant-back interval between treatment and replanting for any crop not listed in the label for this product.
- Do not feed treated vines and hay to livestock.
- Do not treat cowpeas, since these are considered to be grown as livestock feed.

®Trademark of Dow AgroSciences LLC  
R353-014  
EPA accepted 05/12/08  
Replaces R353-007