

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Durango<sup>®</sup> DMA<sup>®</sup>

EPA Reg. No. 62719-556

### Weed Control in Cotton with Roundup Ready<sup>®</sup> Flex Gene in Arizona

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the General Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, Application Equipment and Application Methods, and Roundup Ready Crops on the label attached to this product's container.

See the Roundup Ready Crops section of the product label for general precautionary instructions for use in Roundup Ready Crops.

The use directions in this section apply only to varieties marked as Roundup Ready Flex cotton. If this product is applied over the top of cotton other than Roundup Ready Flex cotton, crop injury and reduced yields will result. Do not combine the directions in this section with the directions in the section for Cotton with Roundup Ready Gene or with any other Roundup Ready cotton or Roundup Ready Flex cotton use directions on other glyphosate-containing products. If this product drifts onto adjacent fields of post 4-leaf (node) Roundup Ready cotton, extensive crop injury, including boll loss, delayed maturity and/or yield loss will occur.

**Types of Applications:** Preplant, at-planting, preemergence, postemergence, preharvest

#### Maximum Allowable Application Rates

Application Type	Rate (quart/acre)
preplant preemergence at-planting	3.75
total all in-crop applications from cracking to 60 percent open bolls	4.25
maximum allowed from 60 percent open bolls to 7 days prior to harvest	1.5
combined total for all applications	6

Calculate the combined rate used for all preplant, in-crop and preharvest applications to ensure that the total does not exceed the maximum rate per acre per year allowed.

### **Preplant, Preemergence, and At-Planting**

Apply this product before, during, or after planting Roundup Ready Flex cotton.

### **Postemergence**

Apply this product to Roundup Ready Flex cotton to control annual grasses and broadleaf weeds listed on this label. Eliminate competing weeds early to maximize yield potential. One or more applications of this product will control or suppress many perennial weeds. For best results, initially apply 24 fl oz per acre on 1- to 3-inch tall annual grass and broadleaf weeds. Apply this product at up to 48 fl oz per acre per application using ground application equipment. More thorough weed coverage may be achieved by using post-directed spray equipment.

### **Precautions and Restrictions:**

- Do not apply more than 48 fl oz of this product per acre as a maximum single in-crop application rate using ground equipment.
- Do not apply more than 24 fl oz per acre as an in-crop application made alone or with the addition of other crop chemicals containing surfactants as a crop response including leaf speckling or leaf necrosis may occur.
- Do not apply more than 36 fl oz of this product per acre when applying by air.
- The maximum combined total application rate of this product between layby and 60 percent open bolls is 48 fl oz per acre.
- The maximum combined total application rate of this product made from crop emergence to 60 percent open bolls is 4.25 quarts per acre.
- Do not add additional surfactant or additives containing surfactant to this product for over the top application to Roundup Ready Flex cotton.

### **Preharvest**

Apply this product to Roundup Ready Flex cotton at 48 fl oz per acre for annual and perennial weed control prior to harvest after 60 percent boll crack. Make the application by aerial or ground spray equipment. When applied to Roundup Ready Flex cotton, this product does not enhance the performance of harvest aids.

### **Precautions and Restrictions:**

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not add additional surfactant or additives containing surfactant to this product for preharvest application to Roundup Ready Flex cotton.

**Attention:** Normal growth of Roundup Ready Flex cotton will result if this product is applied according to label directions. It is impossible to eliminate all risks associated with this product due to the sensitivity of cotton fruiting to various environmental conditions, agronomic practices and other factors even when applications are made in conformance with label specifications. These factors, in some cases, can result in boll loss, delayed maturity and/or yield loss.

®Trademark of Dow AgroSciences LLC  
Roundup Ready® is a registered trademark of Monsanto Company

R353-029  
EPA accepted 11/26/10  
Initial printing