

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Enable[®] 2F

EPA Reg. No. 62719-416

Disease Control in Citrus Fruits[†]

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

[†]Including, but not limited to: Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Orange (Sweet, Sour and Mandarin), Pummello, and Tangerine

Refer to product label for Enable 2F for Mixing and Application Instructions.

Apply aerially in a recommended minimum of 10 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Enable 2F with a nonionic surfactant or spray oil at the manufacturer's recommended rate and making thorough coverage applications.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
greasy spot (<i>Mycosphaerella citri</i>)	8 (0.12 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application or for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> • Do not make more than 3 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year. • Preharvest Interval: Up to day of harvest. • Minimum Re-Treatment Interval: 21 days. • Do not allow livestock to graze in treated areas. • Do not apply Enable 2F with polymer based spray adjuvants.
scab (<i>Elsinoe fawcettii</i>)		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold (<i>Capnodium</i> spp.)		Applications made for control of greasy spot or scab will also control sooty mold.	

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R833-018

EPA accepted: 01/19/07

Initial printing.