

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Enable[®] 2F

EPA Reg. No. 62719-416

Disease Control in Sugarbeets

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Enable[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Enable 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Refer to product label for Enable 2F for Mixing and Application Instructions.

Disease	Enable 2F (fl oz/acre)	Use Directions	Restrictions
cercospora leaf spot (<i>Cercospora beticola</i>) powdery mildew (<i>Erysiphe</i> spp.)	8 (0.125 lb active)	Apply Enable 2F at intervals of 14 to 21 days beginning when conditions are favorable for the development of disease. For resistance management, alternate applications of Enable 2F with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant such as crop oil concentrate that is labeled for crop use should be added to the spray mixture at the manufacturer's recommended rate. Add an adjuvant to the spray solution last. For best spray coverage of foliage, apply aurally in a recommended minimum of 5 to 7 gallons of water per acre or by ground in a recommended minimum of 20 gallons of water per acre.	<ul style="list-style-type: none">• Do not make more than 2 applications or apply more than 16 fl oz of Enable 2F (0.25 lb active) per acre per year.• Preharvest Interval: Do not apply within 14 days of harvest.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, the use of Enable 2F should be part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.2 to 1 lb active per acre, the PBI must be 210 days.

®Trademark of Dow AgroSciences LLC

R833-022

EPA accepted: 01/19/07

Initial printing.