

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Entrust[®]

EPA Reg. No. 62719-282

Foliar Insect Control in Pomegranate

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Entrust[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Entrust according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Entrust.

Directions for Use

Refer to product label for Entrust for General Use Precautions, Mixing, and Application information.

Pests and Application Rates:

Pests	Entrust	
	(oz/acre)	Dilute Spray (oz/100 gal)
carob moth filbert moth leafrollers (such as oblique-banded omnivorous fruit tree pandemis redbanded variegated) light brown apple moth navel orangeworm oriental fruit moth peach twig borer thrips ¹ western cherry fruit fly	1.25 – 2.5	0.42 – 0.83

¹Control of thrips may be improved by addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing.

Application Timing: Optimal timing for **leafrollers** may vary between species and geographic location. Monitor the moth flights and the infestation densities of the larval stages. Thorough coverage is necessary for optimal control. Consult with your Dow AgroSciences representative, state agricultural experiment station, certified pest control advisor or extension specialist for specific application timings in your area.

Application Rate: Choose a higher rate in the rate range for large trees, heavy infestations, or advanced growth stages of target pest, especially if spray volume or coverage is marginal.

Spray Volume: Dilute sprays are sprayed to the point of runoff. The application rate range in the table is based on a spray volume of 300 gallons per acre. Gallonage of dilute sprays will vary depending on tree size, density of canopy, stage of seasonal growth, and spacing in the orchard.

Resistance Management: Do not make more than 3 consecutive applications of Group 5 insecticides (spinetoram and spinosad) within a crop season. If additional treatments are required after 3 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid treating consecutive generations of oriental fruit moth and leafrollers. Do not apply more than 3 sprays targeted at leafrollers per season.

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Minimum Treatment Interval:** A 10- to 14-day re-treatment schedule may be necessary to maintain control if the crop is growing rapidly or if there is heavy pest pressure.
- Do not apply more than a total of 9 oz of Entrust (0.45 lb ai spinosad) per acre per year.
- Do not apply more than 3 sprays targeted at leafrollers per season.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R184-042

EPA accepted: 02/03/10

Replaces R184-033