

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Entrust<sup>®</sup>

EPA Reg. No. 62719-282

### Foliar Insect Control or Suppression in Bulb Vegetables (Crop Group 3<sup>†</sup>)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Entrust<sup>®</sup> Naturalyte<sup>®</sup> insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Entrust according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Entrust.

#### Directions for Use

<sup>†</sup>Bulb vegetables (Crop Group 3) including, but not limited to: Dry Bulb Onion, Garlic, Great-Headed (Elephant) Garlic, Green Onion, Leek, Shallot, Welch Onion

Refer to product label for Entrust for General Use Precautions, Mixing and Application instructions.

#### Pests and Application Rates:

Pests	Entrust (oz/acre)	Specific Use Restriction
armyworms dipteran leafminers European corn borer fleabeetle loopers	1 - 2 (0.05 – 0.1 lb ai/acre)	<b>Do not</b> apply more than a total of 9 oz of Entrust (0.45 lb ai of spinosad) per acre per crop or make more than 5 applications per calendar year.
thrips (suppression) <sup>†</sup>	1.25 – 2.5 (0.06 – 0.12 lb ai/acre)	

<sup>†</sup>Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing on the product label. If thorough coverage is desired, then high pressure (>70 psi) directed sprays with dual directed nozzles can assist leaf penetration of onion.

#### Specific Use Directions:

**Application Timing:** For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Apply as a foliar spray at the rate indicated for target pest. Use a higher rate in specified rate range for larger larvae or heavier infestations.

#### General Use Restrictions:

- **Minimum Re-treatment Interval:** Do not make applications less than 4 days apart.
- **Pre-harvest Interval:** Do not apply within 1 day of harvest.

**Note:** This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R184-008

EPA-accepted 02/24/06

Initial printing.