



FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

For use in lake restoration projects in dewatered zones of lakes

Escort XP Herbicide®

EPA Reg. No. 432-1549

EPA SLN. No. FL-170004

THIS LABEL IS VALID UNTIL 12/31/2021 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELLED, OR SUSPENDED.

Active Ingredient:

Metsulfuron methyl	60%
Other ingredients	40%

Escort XP Herbicide is a dispersible granule that is mixed in water and applied as a spray. The use of Escort XP Herbicide under this Special Local Need registration is specific to application(s) for control of broadleaf weeds in dewatered zones of lakes. Escort XP Herbicide use in lake restoration projects in dewatered zones of lakes is limited to the Florida Fish and Wildlife Conservation Commission, or their contractors.

ENDANGERED SPECIES

Do not apply to areas where the Federally endangered Okeechobee gourd (*Cucurbita okeechobeensis* ssp. *Okeechobeensis*) is present. To determine if the gourd is in the treatment area, which in South Florida is within one mile of Lake Okeechobee levees, contact Florida’s Endangered Species Protection Coordinator at (850) 487-0532.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

Target Pests

Broadleaf weeds, including Pickerelweed (*Pontederia*), Arrowhead (*Sagittaria*), Smartweed (*Polygonum*), and Willow (*Salix*).

Applications to Sites

Lake restoration projects in dewatered zones of lakes.

Application rate:

Apply 0.5-2.0 ounces of Escort XP for broadcast or spot applications. Select a spray volume and delivery system that will ensure thorough coverage and uniform spray pattern.

Do not apply more than 2.0 ounces of Escort XP Herbicide via broadcast application(s), per acre per year. (i.e., every 12 months)

Method of Application:

Broadcast applications may be made by ground application or by aerial application (helicopter only). Spot applications may be made with high pressure sprayer and hand gun, or portable equipment such as backpack sprayers, hand-held sprayers, or spray bottles. A non-ionic surfactant may be used per surfactant label directions.

PRECAUTIONS

- Do not apply this product through any type of irrigation system.
- Do not apply to greater than 10% of the total lake area. [For example, you may treat up to 500 acres of a 5,000 acre lake (normal pool) although 1,000 acres may be exposed after the drawdown.]
- Do not apply within 30 days of refilling the lake or closing the structures on a lake for refilling.
- Do not apply within 50 feet of lake/reservoir water's surface.
- Application should be made only to areas where the receded waters are not used for irrigation or drinking water.

There are no grazing restrictions when Escort XP Herbicide is applied at rates that do not exceed 2 ounces per acre per year.

Refer to the Federal Section 3 label for additional product information, precautions and restrictions.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any pesticide product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer Environmental Science
A Division of Bayer CropScience LP
5000 CentreGreen Way, Suite 400
Cary, NC 27513**