



Bayer Environmental Science, a division of
 Bayer CropScience, LP
 5000 CentreGreen Way, Suite 400
 Cary, NC 27513

ESPLANADE® F

For use on conifer production areas where
 Christmas trees are grown.

EPA Reg. No. 432-1517

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF Michigan, North Carolina, Oregon, Pennsylvania, Washington and Wisconsin

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Esplanade F may be applied for pre-emergent weed control in commercial field grown Christmas tree production areas. Apply as a directed spray to the soil surrounding established plants. Apply 3 - 6 fl oz of Esplanade F per acre for these uses. Applications must only be made after rainfall has settled the soil around the roots following transplanting. Cultivation or disturbance of the soil surface after application is not recommended since it may reduce weed control. Christmas trees that are tolerant to Esplanade F are listed in the table below. Other species may be treated if the user has prior experience with Esplanade F. Without prior experience, it is recommended that small areas be treated with Esplanade F to determine selectivity on specific tree species before large scale applications are made. The user accepts all responsibility for injury on any tree species not listed on this label.

Common Name	Scientific Name	Cultivar
Eastern Redcedar	<i>Juniperus virginiana</i>	Burkii, Idyllwild
Douglas-fir	<i>Pseudotsuga menziesii</i>	
Fir, Fraser	<i>Abies fraseri</i>	
Pine, Scotch (Scots pine)	<i>Pinus sylvestris</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, Eastern White	<i>Pinus strobus</i>	
Spruce, Colorado Blue (Blue spruce)	<i>Picea pungens</i>	
Spruce, Blackhills (White spruce)	<i>Picea glauca</i>	Densata
Spruce, Dwarf Alberta (White spruce)	<i>Picea glauca</i>	Conica
Spruce, Norway	<i>Picea abies</i>	

- Do not exceed 10 fl oz of Esplanade F per acre in a 12-month period
- Do not use in production ornamentals, nursery or greenhouse production sites, hardscapes or landscape ornamentals
- Do not use on trees grown in containers
- Do not apply irrigation to treated areas within 48 hours after application
- Do not apply over-the-top of trees or allow sprays to contact foliage or injury may occur

Soil conditions can affect the tolerance of Christmas trees to Esplanade F. Excessively coarse or sandy soils may allow for downward movement of Esplanade F into the root zone and cause significant root damage and phytotoxicity. Coarse soils may include significant quantities of sand or gravel. Prior to application of Esplanade F on these soils, confirm soil texture with a soil test. Christmas trees grown in soil exceeding 90% sand or 20% gravel may be at risk. If Esplanade F is to be applied in these soils, evaluate tolerance of a few plants of each species in Esplanade F treated soil for 1-2 months prior to a large scale application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).