

TECHNICAL INFORMATION BULLETIN

GROUP 2 HERBICIDE

Everest[®] 3.0 *herbicide*

EPA Reg. No. 66330-429

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF:
MONTANA, NORTH DAKOTA and SOUTH DAKOTA**

Tank-Mix of EVEREST[®] 3.0 with Talinor[®]¹

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered labels for **EVEREST[®] 3.0 (EPA Reg. No. 66330-429)** and **Talinor[®] (EPA Reg. No. 100-1570)**.

¹ Talinor herbicide contains the active ingredients bicyclopyrone and bromoxynil

120816V005MTNDS2ee

A tank-mix of EVEREST 3.0 at (1.5 - 2.0 fl oz/A), Talinor at (13.7 - 18.2 fl oz/A) and CoAct+™ spray additive at (2.75 - 3.6 fl oz/A) can be used in a postemergence application to improve control of broadleaf weeds listed on the EVEREST 3.0 label. This tank-mixture can be applied to spring, winter and durum wheat from 2-leaf stage up to 60 days prior to harvest.

USE RATES

EVEREST 3.0 Rate (fl oz/A) ²	Talinor Rate (fl oz/A)	CoAct+ Rate (fl oz/A)	Non-Ionic Surfactant (% v/v)
1.5 – 2.0	13.7	2.75	0.25
1.5 – 2.0	16	3.2	0.25
1.5 – 2.0	18.2	3.6	0.25

² EVEREST 3.0 can be used at a reduced rate in spring wheat, including durum wheat, for high pH and light texture soils. 1.5 fl oz/A can be used when soil pH is 7.8 or greater and if organic matter is less than 3%.

RESTRICTIONS:

- Follow all applicable directions, restrictions and precautions on the labels of all products used in the tank-mix. This includes mixing instructions.
- Apply only to spring, winter and durum wheat. Do not apply to barley.
- Always add the directed rate of CoAct+ to the spray solution.
- Do not add ammonium sulfate (AMS) containing products to the spray mixture as severe crop injury may occur.
- Always use a non-ionic surfactant (NIS) containing at least 80% active ingredient at 0.25% v/v (1 quart/100 gallons) finished spray volume.
- Do not mix with organophosphate or carbamate insecticides as severe crop injury may occur
- Do not apply organophosphate or carbamate insecticides within 7 days before 7 days after application of this tank mixture. Severe crop injury may occur.

Produced For:

ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, North Carolina 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

EVEREST is a registered trademark of an Arysta LifeScience Group Company.

Talinor is a registered trademark of a Syngenta Company.

CoAct+ is a trademark of Syngenta Participations AG.

©2018 Arysta LifeScience Group Company.